



Maternal mortality and morbidity burden in the Eastern Mediterranean Region: findings from the Global Burden of Disease 2015 study

GBD 2015 Eastern Mediterranean Region Maternal Mortality Collaborators ·
Ali H. Mokdad¹

Received: 1 May 2017/Revised: 15 June 2017/Accepted: 21 June 2017
© The Author(s) 2017. This article is an open access publication

Abstract

Objectives Assessing the burden of maternal mortality is important for tracking progress and identifying public health gaps. This paper provides an overview of the burden of maternal mortality in the Eastern Mediterranean Region (EMR) by underlying cause and age from 1990 to 2015.

Methods We used the results of the Global Burden of Disease 2015 study to explore maternal mortality in the EMR countries.

Results The maternal mortality ratio in the EMR decreased 16.3% from 283 (241–328) maternal deaths per 100,000 live births in 1990 to 237 (188–293) in 2015. Maternal mortality ratio was strongly correlated with socio-demographic status, where the lowest-income countries contributed the most to the burden of maternal mortality in the region.

Conclusion Progress in reducing maternal mortality in the EMR has accelerated in the past 15 years, but the burden remains high. Coordinated and rigorous efforts are needed to make sure that adequate and timely services and

interventions are available for women at each stage of reproductive life.

Keywords Maternal mortality · Maternal health · Eastern Mediterranean Region · Burden of disease

Introduction

Maternal mortality ratio (MMR), which measures deaths per 100,000 live births, is a standard measure for global, regional, and national comparison (Abouzahir and Wardlaw 2001). It is also one of the main criteria of health outcomes and an indicator of the socioeconomic development level of countries that is recognized worldwide (Liang et al. 2010). Country estimates of maternal mortality over time are crucial to inform the planning of maternal, sexual, and reproductive health programs and to guide advocacy efforts and research at the national level. These estimates are also needed at the international level to inform decision-making concerning resource allocation by development partners and donors (WHO et al. 2007).

In 2015, 275,288 women are estimated to have died as a result of pregnancy or childbirth globally (Kassebaum et al. 2016b). Maternal death is defined by ICD-10 as the death of a woman while pregnant or within 42 days and up to 1 year (late maternal death); of termination of pregnancy, irrespective of the duration and site of the pregnancy, from any cause related to or aggravated by the pregnancy or its management, but not from accidental or incidental causes (World Health Organization 1992).

The primary causes of maternal deaths are hemorrhage, hypertension, infections, pre-existing medical conditions, and lack of literacy, family planning (unmet need and birth spacing), and access to pregnancy termination when

Corresponding author: Ali H. Mokdad

This article is part of the supplement “The state of health in the Eastern Mediterranean Region, 1990–2015”.

GBD 2015 Eastern Mediterranean Region Maternal Mortality Collaborators are listed at the end of the article.

Electronic supplementary material The online version of this article (doi:10.1007/s00038-017-1004-3) contains supplementary material, which is available to authorized users.

✉ Ali H. Mokdad
mokdaa@uw.edu

¹ Institute for Health Metrics and Evaluation, University of Washington, Seattle, WA, USA

needed. Identifying and improving the main factors related to maternal mortality depend on a correct definition of the required priorities for appropriate prevention, diagnosis, and treatment (Ribeiro et al. 2008).

Poverty, illiteracy, malnutrition, and the low social status of women are usually among the major underlying causes of maternal mortality. Analyzing the economic, social, and health system factors affecting maternal mortality can provide some credible information and evidence for public health interventions to improve maternal health. A few general studies on nonclinical determinants of maternal mortality have analyzed different factors including access to safe drinking water, access to food, fertility rate, education level, life expectancy at birth, access to health services, delivery done by skilled birth attendants, out-of-pocket payment, GDP per capita, health expenditures, and government corruption (Hertz et al. 1994; Midhet et al. 1998; Buor and Bream 2004; Alvarez et al. 2009; Muldoon et al. 2011).

From 1990 to 2015, the global MMR decreased 30.6% from 280.4 (95% uncertainty interval (UI): 262.6–299.0) to 194.7 (172.6–233.2) (Institute for Health Metrics and Evaluation 2017). This decrease was driven in part by the recognition of maternal health as one of the priorities in the UN Millennium Development Goals (MDGs), which provided a strong incentive to reduce MMR by country (Ronsmans and Graham 2006). New goals outlined in the sustainable development goals (SDGs) for 2015 to 2030 build on the momentum of the MDGs and seek to bring the global MMR below 70 deaths per 100,000 live births (United Nations 2015). The greatest challenge in meeting this goal lies with countries in the bottom two quintiles of wealth, where improvement is most needed and has historically occurred slowly if at all (Abouzahir and Wardlaw 2001).

The variation in MMR between countries by income group is drastic, where the burden of maternal mortality falls almost exclusively upon developing countries (Ronsmans and Graham 2006). Globally, the average MMR for countries in the top quintile of wealth for 2015 was 14.5 (13.7–15.4) compared to 440.2 (357.4–543.5) in the bottom quintile (Institute for Health Metrics and Evaluation 2017). Despite this variation, inequalities in maternal mortality exist between and within countries, where mortality in all locations is exacerbated by a lack of access to quality obstetric care, whether because of the remoteness of certain regions or the inability of health facilities to provide proper care (Peterson et al. 2012).

The Eastern Mediterranean Region (EMR) is a diverse region consisting of 22 countries, including high-income countries, where socioeconomic development has progressed considerably over the last decades; middle-income countries that have well-developed health service delivery

infrastructures but face resource constraints; and low-income countries that lack the resources and infrastructure for effective health interventions. Many countries in the region are suffering from political instability, conflicts, and other complex development challenges. This range from low to high incomes can be seen as a microcosm of the variability in maternal mortality present globally. This paper seeks to examine in detail the burden of maternal mortality within the EMR using the results of the GBD 2015 study.

Methods

A detailed methodology of maternal mortality estimation for GBD 2015 has been published elsewhere (Kassebaum et al. 2016b). In short, sources with population-level data were processed using standardized algorithms to adjust for age-specific, year-specific, and geography-specific patterns of incompleteness. Additional steps were taken to account for underestimations of maternal mortality in vital registration systems. ICD-10 vital registration codes pertaining to HIV-related maternal deaths were excluded and quantified separately (Kassebaum et al. 2014). Overall maternal mortality was modeled using cause-of-death ensemble modeling (CODEm), where all combinations of covariates were tested and ranked on the basis of out-of-sample predictive validity performance. All data sources are available on our website, and we provide a visualization to show each source used in our analyses (Institute for Health Metrics and Evaluation 2016).

We calculated years of life lost (YLLs) by multiplying deaths by the residual expected individual lifespan at the age of death as derived from the GBD 2015 standard model life table. Years lived with disability (YLDs) were calculated by multiplying the number of prevalent cases of a certain health outcome by the disability weight assigned to this health outcome. A disability weight reflects the magnitude of the health loss associated with an outcome and has a value that is anchored between 0, equivalent to full health, and 1, equivalent to death. YLLs were calculated by multiplying deaths by the remaining life expectancy at the age of death from a standard life table chosen as the norm for estimating premature mortality in GBD. Disability-adjusted life years (DALYs) were calculated by adding YLLs and YLDs. Detailed methods on YLLs, YLDs, and DALYs are published elsewhere (Kassebaum et al. 2016a; Vos et al. 2016; Wang et al. 2016).

We evaluated the associations between maternal mortality and socio-demographic status using the socio-demographic index (SDI). SDI is a composite measure developed for GBD 2015 that accounts for fertility rate, lag-dependent income per capita, and education. To

capture the average relationships for each age group, we applied a simple least squares spline regression of the maternal mortality rate on SDI. The SDI is scaled from 0 to 1, where 0 represents the lowest possible observed SDI and 1 is the highest. We reported uncertainty for all our estimates (Kassebaum et al. 2016a).

Results

Mortality

The total number of deaths due to maternal disorders increased 4.5% from 38,595 (32,859–44,785) deaths in 1990 to 40,338 (31,965–49,954) in 2015 in the EMR, compared to a 29.5% decrease globally (Institute for Health Metrics and Evaluation 2017). In 1990, the maternal mortality ratio was similar in the EMR and globally at 283 (241–328) maternal deaths per 100,000 live births in the EMR and 280 (263–299) globally (Institute for Health Metrics and Evaluation 2017). By 1995, a gap between the regional and global trends had opened, and this persisted steadily through 2015. By then, the EMR had a maternal mortality ratio of 237 (188–293) compared to 194 (173–223) globally (Institute for Health Metrics and Evaluation 2017). By country, only Afghanistan and Djibouti had an annual increase in maternal mortality ratio between 1990 and 2015 (Table 1). Palestine also had an increase in maternal mortality ratio between 2000 and 2015, but still had a lower ratio in 2015 compared to 1990 (Table 1).

Both globally and regionally, maternal hemorrhage was the leading cause of maternal mortality in 1990 and 2015 (Institute for Health Metrics and Evaluation 2017). By country within the EMR, hemorrhage was also the most common, followed by other maternal disorders, noticeably within the Gulf states (Fig. 1). In all countries, all causes of maternal mortality decreased from 1990 to 2015 (Table 2). In 2015, maternal hemorrhage in Somalia was the largest contributor to maternal mortality by country and causes at 72.3 (13.2–229.0) deaths per 100,000 among women aged 15–49 (Table 2). The GBD study expanded the analysis to the full reproductive age range of 10–54 years (Kassebaum et al. 2016b). In general, the countries with the highest mortality rates by cause also showed the least improvement from 1990 to 2015 (Table 2).

YLLs

YLL rates fell 49% in the EMR between 1990 and 2015, the same as the global mean, from 2,683 (2284–3103) per 100,000 women aged 15–49 to 1377 (1091–1695) (Table 3). In that time frame, the YLL rate decreased in all countries (Table 3). There was significant inter-country

variation in YLL rates. The highest YLL rate in 2015 was in Somalia with 7774 (1497–23,408) YLLs per 100,000 women aged 15–49, compared to Kuwait at 21 (16–27) (Table 3). Somalia also had the smallest percent change from 1990 to 2015, a decrease of 18% (Table 3).

YLDs

Average YLD rates in the region fell from 161 (114–215) per 100,000 women aged 15–49 to 82 (57–113), a 49% decrease (Table 3). Globally, there was a 41% decrease (Institute for Health Metrics and Evaluation 2017). Inter-country variability in the YLD rate was less than that of YLLs, with the highest seen in Somalia with 254 (171–362) YLDs per 100,000 women aged 15–49 and the lowest in the United Arab Emirates with 16 (10–26) (Table 3).

YLL/YLD

The YLL/YLD ratio was included as an indicator of health system effectiveness in dealing with fatal and non-fatal outcomes of maternal disorders. Large values, such as that of Afghanistan, a ratio of 89.6 in 2015, demonstrated cases, where fatal outcomes heavily outweigh non-fatal ones (Table 3). Kuwait in 2015 had a ratio of 1.11, signifying a near equal burden of fatal and non-fatal outcomes for maternal disorders (Table 3). Lower income countries with SDI less than 0.5 all had ratios greater than 10, whereas higher income countries with SDI greater than 0.7 had ratios under 4 (Table 3). In the countries with SDI less than 0.5, all but Sudan had an increase in YLL/YLD ratio from 1990 to 2015 (Table 3). Large decreases in the ratio from 1990 to 2015 highlighted effective improvements in treatment, such as in Morocco, where the ratio decreased from 35.2 to 8.5 (Table 3).

DALYs

From 1990 to 2015, DALY rates decreased in all countries in the region. The overall DALY rate for the region decreased by 49% during this interval, compared to 51% globally (Institute for Health Metrics and Evaluation 2017). As discussed in the YLL/YLD section above, the proportional contribution of YLDs to the DALYs rate increased as SDI increased (Table 3). Like YLL rates, DALY rates showed large inter-country variation, as shown in Fig. 2. DALY rates peaked in the 20–25-year age group, decreasing until ages 35–40, then steeply falling off in 5-year increments (Fig. 3). By sub-cause, complication of abortions contributed the most to the DALY rate in young (10–15) and old (45+) age groups (e-Fig. 1). Percent of DALYs attributable to obstructed labor increased steadily with age, spiking drastically for the 50–55-year age group (e-Fig. 1).

Table 1 Maternal deaths, maternal mortality ratio, and annualized change in maternal mortality ratio in the Eastern Mediterranean Region, 1990–2015 (Global Burden of Disease 2015 study, Eastern Mediterranean countries, 1990–2015)

Country	Number of maternal deaths			Maternal mortality ratio (per 100,000 livebirths)			Annualized rate of change in maternal mortality ratio (%)		
	1990	2000	2015	1990	2000	2015	1990–2000	2000–15	1990–2015
Afghanistan	4590 (2825–7111)	7328 (4529–11,007)	8525 (5010–13,221)	732.3 (451.0–1130.9)	753.3 (466.5–1130.7)	788.9 (464.1–1219.2)	0.4 (–4.8–5.0)	0.3 (–3.6–3.8)	0.3 (–2.1–2.7)
Bahrain	8 (6–11)	5 (4–7)	5 (4–7)	53.6 (39.9–73.5)	35.6 (27.3–47.3)	24.6 (18.3–35.4)	–4.0 (–7.9 to –0.6)	–2.5 (–5.1 to 0.4)	–3.1 (–4.9 to –1.3)
Djibouti	86 (35–159)	118 (43–283)	107 (43–299)	378.3 (154.4–698.7)	523.8 (192.3–1247.0)	486.2 (193.8–1353.5)	2.8 (–9.8 to 15.3)	–0.5 (–9.4 to 9.9)	0.6 (–4.1 to 6.6)
Egypt	2744 (2308–3176)	1186 (976–1414)	1052 (809–1340)	146.7 (123.4–169.7)	69.2 (57.0–82.5)	42.3 (32.5–53.9)	–7.5 (–9.7 to –5.5)	–3.3 (–5.3 to –1.2)	–5.0 (–6.1 to –3.7)
Iran	1039 (704–1432)	426 (337–539)	281 (192–421)	56.6 (38.5–78.0)	34.5 (27.3–43.7)	20.8 (14.2–31.1)	–5.0 (–8.6 to –0.6)	–3.5 (–6.4 to –0.2)	–4.1 (–6.0 to –1.9)
Iraq	969 (702–1300)	950 (655–1311)	729 (428–1199)	146.0 (105.9–195.9)	113.4 (78.3–156.4)	58.6 (34.4–96.1)	–2.5 (–6.7 to 1.4)	–4.5 (–8.2 to –1.0)	–3.7 (–6.0 to –1.3)
Jordan	116 (89–149)	121 (88–159)	48 (36–63)	97.9 (74.6–125.4)	81.2 (58.7–106.2)	24.2 (17.9–31.5)	–1.9 (–5.1 to 1.1)	–8.0 (–10.7 to –5.4)	–5.6 (–7.2 to –4.1)
Kuwait	4 (3–5)	5 (4–6)	4 (3–6)	9.7 (8.1–11.8)	11.9 (10.4–13.5)	5.9 (4.4–7.5)	2.1 (0.0–4.1)	–4.8 (–6.8 to –2.8)	–2.0 (–3.4 to –0.7)
Lebanon	24 (15–38)	17 (11–24)	13 (8–21)	35.6 (22.8–55.1)	26.7 (18.0–39.0)	15.3 (9.0–24.0)	–2.9 (–7.6 to 1.6)	–3.9 (–7.6 to 0.3)	–3.3 (–6.2 to –0.7)
Libya	34 (24–45)	28 (20–37)	30 (20–43)	26.6 (19.1–36.0)	23.9 (17.2–31.5)	22.8 (15.6–32.9)	–1.0 (–4.8 to 2.3)	–0.3 (–3.5 to 2.8)	–0.6 (–2.6 to 1.3)
Morocco	2441 (1891–3201)	1197 (894–1560)	479 (299–756)	332.7 (258.7–436.1)	192.0 (143.4–249.6)	68.5 (42.8–108.1)	–5.4 (–8.8 to –2.4)	–7.0 (–10.1 to –3.4)	–6.4 (–8.6 to –4.3)
Pakistan	16,973 (13,060–21,189)	22,038 (17,789–26,598)	19,005 (14,012–24,369)	391.5 (301.3–488.4)	498.9 (404.1–601.7)	348.6 (257.2–447.0)	2.4 (–0.3 to 5.5)	–2.4 (–4.7 to –0.3)	–0.5 (–1.9 to 0.9)
Palestine	29 (19–43)	17 (14–21)	25 (17–35)	29.3 (19.0–43.7)	14.2 (11.3–17.8)	16.2 (11.5–23.2)	–7.1 (–11.2 to –3.1)	0.8 (–1.9 to 3.9)	–2.4 (–4.7 to 0.2)
Oman	24 (14–38)	12 (8–18)	13 (9–18)	34.1 (20.0–54.8)	21.4 (14.8–31.7)	15.5 (11.0–21.6)	–4.5 (–8.7 to –0.3)	–2.2 (–5.5 to 1.2)	–3.1 (–5.6 to –0.7)
Qatar	7 (4–10)	7 (5–10)	7 (4–9)	64.0 (42.0–91.5)	59.9 (41.4–84.2)	25.5 (16.4–35.7)	–0.6 (–4.6 to 3.6)	–5.7 (–9.5 to –2.4)	–3.7 (–6.0 to –1.6)
Saudi Arabia	124 (93–162)	101 (86–119)	97 (82–115)	21.3 (16.0–27.9)	17.9 (15.2–21.0)	15.7 (13.3–18.5)	–1.6 (–4.0 to 0.6)	–0.9 (–2.4 to 0.6)	–1.2 (–2.5 to 0.1)
Somalia	2509 (418–6 772)	2899 (489–7449)	3443 (661–10 512)	830.5 (138.5–2193.9)	811.0 (137.5–2077.9)	731.1 (140.5–2218.6)	–0.3 (–11.8 to 10.4)	–0.8 (–8.6 to 8.8)	–0.6 (–6.7 to 6.5)

Table 1 continued

Country	Number of maternal deaths				Maternal mortality ratio (per 100,000 livebirths)				Annualized rate of change in maternal mortality ratio (%)			
	1990	2000	2015	1990	2000	2015	1990-2000	2000-15	1990-2000	2000-15	1990-2015	
Sudan	4131 (2736-5570)	4653 (2790-6731)	3941 (2098-5999)	485.5 (321.7-654.4)	414.4 (248.7-598.0)	298.8 (159.2-454.7)	-1.5 (-5.7-1.9)	-2.3 (-5.5 to 1.0)	-1.5 (-5.7-1.9)	-2.3 (-5.5 to 1.0)	-2.0 (-4.3 to 0.1)	
Syria	560 (421-719)	365 (276-480)	237 (181-309)	125.7 (94.6-161.6)	73.6 (55.6-96.7)	54.1 (41.3-70.5)	-5.3 (-2.6)	-2.0 (0.5)	-5.3 (-8.1 to -2.6)	-2.0 (-4.7 to 0.5)	-3.4 (-5.0 to -1.8)	
Tunisia	183 (143-231)	99 (74-126)	82 (54-115)	85.6 (66.9-108.0)	59.2 (44.3-75.2)	40.6 (26.6-56.9)	-3.7 (-0.8)	-2.6 (0.2)	-3.7 (-6.6 to -0.8)	-2.6 (-5.5 to 0.2)	-3.0 (-4.9 to -1.4)	
United Arab Emirates	15 (9-23)	12 (9-17)	18 (11-28)	31.8 (19.9-48.8)	24.0 (17.0-32.4)	18.0 (10.7-28.3)	-2.6 (2.0)	-2.0 (2.0)	-2.6 (-7.3 to 2.0)	-2.0 (-6.1 to 2.0)	-2.3 (-5.3 to 0.4)	
Yemen	2559 (1342-3890)	2810 (1509-4648)	2631 (1402-5009)	402.6 (211.8-611.4)	399.6 (215.7-659.8)	307.4 (164.5-583.8)	-0.3 (4.5)	-1.9 (2.1)	-0.3 (-4.1 to 4.5)	-1.9 (-5.5 to 2.1)	-1.3 (-3.7 to 1.8)	

Discussion


Our study showed that maternal mortality has declined in the EMR except in Djibouti, Palestine, and Afghanistan in recent years. However, the burden is still higher than the global average. Our findings call for increased efforts to reduce maternal mortality in the region. Moreover, with unrest in the region, there is a risk of losing some of the observed gains.

Efforts to reduce maternal mortality in the EMR have varied in scope, approach, and success by country. This has resulted in great variations and disparities in maternal mortality levels between countries. The high-income countries in the region (Kuwait, Oman, Qatar, Saudi Arabia, and United Arab Emirates) have achieved reductions between 25 and 50% compared to levels in 1990. Maternal mortality ratios in these countries ranged from 6 to 26 deaths per 100,000 live births, consistent with countries of similar income around the world. Certain middle-income countries such as Morocco and Jordan have successfully implemented interventions that have drastically reduced maternal mortality by greater than 75% from 1990 levels. Low-income countries have seen the least improvement, or in some cases none at all. Reduction in maternal mortality in Pakistan, Somalia, Sudan, and Yemen ranged from 11 to 39%, and maternal mortality in Afghanistan and Djibouti increased from 1990 to 2015. Maternal mortality ratios for these countries ranged from 299 deaths per 100,000 live births in Sudan to 789 in Afghanistan. This discussion will examine the obstacles to effective maternal care, the direct and underlying causes of maternal mortality, and the challenges faced by countries in the region to improve maternal health.

The process of improving maternal health in the region faces many challenges which have slowed progress in recent years. For many years, the EMR has been devastated by manmade disasters and conflicts, which have destroyed infrastructure in several nations and have tremendously affected the health of the population, especially for vulnerable groups such as mothers and children. The humanitarian community has made several efforts to address these challenges, and these have reduced the potential impact of the unrest. However, the disruption of health care systems has had a negative impact. The social and health impacts of political instability, domestic crises, and economic sanctions are well-documented in the region (Mokdad et al. 2016). Recent distress in the region will result in deteriorating health conditions in these countries for many years to come. Despite this, however, the region has experienced improved health and life expectancy.

The region is in need of efforts to improve preventive strategies for reducing maternal mortality. A literature

Fig. 1 Top cause of maternal deaths in Eastern Mediterranean Region countries, 2015 (Global Burden of Disease 2015 study, Eastern Mediterranean Countries 2015)



review (Warren et al. 2015) that evaluated the evidence on the effectiveness of sexual and reproductive health interventions delivered in humanitarian crises found some evidence to support increased access and demand creation for family planning services through community health workers, health care subsidies, and discussions within literacy groups. Involving communities in maternal and child health and birth preparedness programs, as well as refurbishing clinics and hospital facilities, was also associated with increased positive health outcomes (Warren et al. 2015).

Female genital mutilation (FGM) is still practiced in many countries in the EMR due to a mix of sociocultural factors within families and communities (UNICEF 2016). Experience of FGM increases the short- and long-term health risks to women and young girls between infancy and age 15, and is unacceptable from a human rights and health perspective. Procedures can cause severe bleeding and problems urinating, and later, cysts, infections, as well as complications in childbirth and increased risk of newborn deaths (WHO 2017).

Thaddeus and Maine provide a conceptual model for understanding factors that obstruct favorable obstetric outcomes that consists of three types of delays a pregnant woman faces when seeking care—delays in deciding to seek care, in reaching an adequate facility, and in receiving adequate care in a facility. The first delay manifests itself in the way women and their families perceive the accessibility of services, often shaped by prior experiences with health facilities. Any number of factors may influence a decision to delay treatment, such as cost, distance, lack of proper medical supplies, and unhelpful or impolite staff. The second delay accounts for the time taken to reach a medical facility. This is a particularly large obstacle for rural areas in which the availability of transportation is uncertain or

expensive. In addition, data on maternal deaths that occur on the way to a health facility are scarce. The third and final delays are in receiving adequate care at health facility, which can be attributed to insufficient staff, drugs, and equipment. It is necessary for interventions to effectively address the underlying causes of these delays on a country-by-country basis (Thaddeus and Maine 1994).

This conceptual model is backed by an extensive body of research on the direct and underlying causes of maternal mortality. Improving women's literacy and knowledge can significantly contribute to their health. Alvarez et al. reported a significant negative relationship between MMR and gross domestic product (GDP) per capita, health expenditure (HE), women's literacy rate, and the number of deliveries by skilled birth attendants in the study region in Africa (Alvarez et al. 2009). Another study in sub-Saharan African countries, concluded that birth in the presence of health professionals and the life expectancy at birth are highly correlated to maternal mortality. Furthermore, a convincing relationship was found between GDP per capita and health expenditure per capita and maternal mortality (Buor and Bream 2004). Similarly, a study in Pakistan reported that a significant relationship was observed between having access to health services, professional health staff, and health care during pregnancy and reductions in MMR (Midhet et al. 1998). The positive impacts of GDP, HE, and female education level on health outcomes were also reported in other studies in the EMR (Bayati et al. 2013; Homaie Rad et al. 2013).

Success stories do exist in the region: for example, Morocco was one of only ten countries globally that met the MDG 5 requirement of an annual rate of decline exceeding 5.5% for maternal mortality ratio every year from 1990 to 2015 (Kassebaum et al. 2016b). This improvement can be traced to a number of programs

Table 2 Mortality rates for causes of maternal mortality in the Eastern Mediterranean Region countries, ages 15–49, 2015 and percent change 1990–2015 (Global Burden of Disease 2015 study, Eastern Mediterranean countries, 1990–2015)

Country	Maternal hemorrhage		Maternal hypertension		Other maternal		Complication of abortion	
	Mortality rate per 100,000, 2015	% change, 1990–2015	Mortality rate per 100,000, 2015	% change, 1990–2015	Mortality rate per 100,000, 2015	% change, 1990–2015	Mortality rate per 100,000, 2015	% change, 1990–2015
Eastern Mediterranean Region	7.61 (5.51–10.37)	-52.27	4.49 (3.29–6.11)	-40.39	3.27 (2.37–4.34)	-48.51	2.59 (1.84–3.53)	-50.05
Afghanistan	47.70 (27.17–77.35)	-36.96	20.12 (11.20–32.87)	-29.85	13.99 (7.81 to 23.27)	-33.95	12.82 (7.07–21.40)	-35.04
Bahrain	0.28 (0.19–0.42)	-80.69	0.31 (0.21–0.47)	-73.93	0.43 (0.30–0.65)	-77.48	0.22 (0.15–0.33)	-74.69
Djibouti	17.36 (6.43–49.10)	-30.30	7.26 (2.58–20.67)	-29.47	4.04 (1.39–11.26)	-30.29	3.61 (1.12–9.89)	-30.47
Egypt	1.90 (1.43–2.48)	-81.48	0.76 (0.54–0.99)	-71.80	0.32 (0.24–0.44)	-55.19	0.59 (0.44–0.78)	-73.29
Iran	0.29 (0.19–0.45)	-88.16	0.17 (0.11–0.27)	-81.40	0.13 (0.09–0.21)	-88.48	0.14 (0.09–0.22)	-88.12
Iraq	3.04 (1.71–5.24)	-70.49	1.04 (0.56–1.85)	-61.10	0.91 (0.47–1.63)	-66.49	1.74 (0.96–2.85)	-66.30
Jordan	0.38 (0.25–0.54)	-89.46	0.34 (0.23–0.49)	-83.99	0.99 (0.69–1.34)	-82.45	0.18 (0.12–0.26)	-86.30
Kuwait	0.07 (0.04–0.09)	-52.16	0.04 (0.03–0.06)	-31.67	0.10 (0.07–0.14)	-48.26	0.09 (0.07–0.13)	-44.51
Lebanon	0.1*6 (0.09–0.27)	-80.68	0.18 (0.10–0.29)	-73.17	0.16 (0.09–0.26)	-76.51	0.10 (0.06–0.17)	-78.00
Libya	0.42 (0.26–0.64)	-54.16	0.26 (0.16–0.41)	-45.63	0.34 (0.21–0.53)	-51.63	0.26 (0.15–0.43)	-46.24
Morocco	1.17 (0.71–1.95)	-89.64	1.19 (0.72–1.91)	-85.15	1.06 (0.64–1.70)	-86.86	0.57 (0.34–0.92)	-88.82
Oman	0.18 (0.11–0.28)	-83.85	0.34 (0.22–0.50)	-77.27	0.35 (0.22–0.52)	-80.07	0.25 (0.16–0.36)	-82.47
Pakistan	9.63 (6.44–13.52)	-52.83	8.81 (5.88–12.43)	-35.15	5.54 (3.46–7.77)	-44.39	4.08 (2.57–6.09)	-47.04
Palestine	0.47 (0.30–0.72)	-70.21	0.38 (0.24–0.58)	-59.52	0.34 (0.22–0.53)	-67.21	0.49 (0.32–0.73)	-65.73
Qatar	0.23 (0.13–0.35)	-83.54	0.35 (0.21–0.52)	-76.08	0.38 (0.23–0.57)	-79.73	0.16 (0.09–0.24)	-80.55
Saudi Arabia	0.15 (0.11–0.20)	-74.35	0.09 (0.06–0.12)	-66.88	0.50 (0.40–0.63)	-59.58	0.25 (0.18–0.33)	-68.39
Somalia	72.34 (13.24–229.05)	-17.78	15.14 (2.95–47.52)	-17.10	10.88 (2.01–33.59)	-18.94	11.63 (2.13–35.48)	-17.53
Sudan	8.52 (4.32–13.68)	-64.24	5.09 (2.55–8.46)	-51.00	5.49 (2.83–9.17)	-55.11	3.02 (1.51–4.90)	-53.80
Syria	1.70 (1.18–2.34)	-77.15	0.66 (0.43–0.97)	-69.42	1.17 (0.82–1.62)	-74.60	0.33 (0.22–0.48)	-72.31
Tunisia	1.00 (0.60–1.46)	-72.70	0.36 (0.22–0.55)	-67.89	0.48 (0.30–0.72)	-68.33	0.20 (0.11–0.32)	-67.14
United Arab Emirates	0.18 (0.10–0.31)	-79.78	0.24 (0.13–0.39)	-73.34	0.27 (0.15–0.46)	-78.40	0.18 (0.10–0.30)	-76.09
Yemen	12.07 (6.03–24.02)	-64.23	5.40 (2.66–10.39)	-55.94	7.08 (3.51–14.63)	-60.55	5.48 (2.70–10.85)	-63.71
Country	Maternal indirect		Obstructed labor		Maternal sepsis		Maternal late	
	Mortality rate per 100,000, 2015	% change, 1990–2015	Mortality rate per 100,000, 2015	% change, 1990–2015	Mortality rate per 100,000, 2015	% change, 1990–2015	Mortality rate per 100,000, 2015	% change, 1990–2015
Eastern Mediterranean Region	2.05 (1.40–2.91)	-48.67	1.93 (1.33–2.65)	-38.25	1.86 (1.32–2.57)	-54.00	0.63 (0.37–1.00)	-46.51
Afghanistan	6.94 (3.71–11.87)	-33.85	6.56 (3.57–10.95)	-28.66	4.49 (2.42–7.52)	-34.04	1.82 (0.90–3.29)	-30.65

Table 2 continued

Country	Maternal indirect		Obstructed labor		Maternal sepsis		Maternal late	
	Mortality rate per 100,000, 2015	% change, 1990–2015	Mortality rate per 100,000, 2015	% change, 1990–2015	Mortality rate per 100,000, 2015	% change, 1990–2015	Mortality rate per 100,000, 2015	% change, 1990–2015
Bahrain	0.18 (0.12–0.27)	-77.82	0.02 (0.01–0.03)	-74.29	0.09 (0.06–0.13)	-76.25	0.03 (0.01–0.04)	-77.72
Djibouti	5.95 (2.00–16.88)	-30.94	1.36 (0.38–4.09)	-31.64	2.51 (0.80–7.26)	-30.79	1.88 (0.55–5.39)	-26.60
Egypt	0.13 (0.07–0.21)	-83.00	0.16 (0.11–0.23)	-61.01	0.55 (0.39–0.74)	-83.76	0.06 (0.04–0.09)	-77.93
Iran	0.24 (0.16–0.37)	-81.40	0.05 (0.03–0.08)	-85.86	0.11 (0.07–0.17)	-85.22	0.02 (0.01–0.04)	-85.56
Iraq	0.42 (0.21–0.76)	-68.17	0.18 (0.09–0.33)	-56.85	0.67 (0.36–1.20)	-65.77	0.15 (0.07–0.28)	-65.80
Jordan	0.19 (0.12–0.28)	-87.75	0.11 (0.07–0.17)	-84.20	0.20 (0.13–0.29)	-86.93	0.04 (0.03–0.07)	-86.91
Kuwait	0.04 (0.03–0.06)	-43.63	0.01 (0.01–0.02)	-36.29	0.04 (0.02–0.05)	-39.62	0.01 (0.00–0.01)	-43.71
Lebanon	0.08 (0.04–0.14)	-76.84	0.03 (0.01–0.06)	-74.99	0.07 (0.04–0.12)	-76.85	0.02 (0.01–0.03)	-76.89
Libya	0.15 (0.09–0.25)	-55.50	0.06 (0.03–0.10)	-42.48	0.16 (0.09–0.25)	-51.42	0.04 (0.02–0.06)	-49.31
Morocco	0.24 (0.13–0.41)	-86.68	0.34 (0.19–0.59)	-82.51	0.45 (0.27–0.74)	-86.78	0.06 (0.03–0.10)	-87.29
Oman	0.17 (0.11–0.26)	-80.48	0.02 (0.01–0.03)	-78.77	0.07 (0.04–0.10)	-81.47	0.02 (0.01–0.04)	-79.45
Pakistan	2.75 (1.69–4.11)	-44.69	4.84 (3.21–6.91)	-37.12	2.88 (1.77–4.28)	-44.67	1.16 (0.62–1.90)	-45.07
Palestine	0.18 (0.11–0.29)	-67.34	0.04 (0.02–0.07)	-62.46	0.12 (0.07–0.19)	-67.65	0.07 (0.04–0.12)	-64.98
Qatar	0.23 (0.14–0.36)	-81.22	0.02 (0.01–0.03)	-78.54	0.22 (0.14–0.35)	-78.62	0.07 (0.04–0.11)	-74.65
Saudi Arabia	0.07 (0.05–0.10)	-73.74	0.02 (0.01–0.03)	-70.76	0.14 (0.10–0.18)	-70.85	0.02 (0.01–0.03)	-69.15
Somalia	15.86 (2.85–50.91)	-19.42	3.35 (0.53–11.26)	-21.48	6.66 (1.16–21.53)	-20.07	5.38 (0.95–17.98)	-17.88
Sudan	7.44 (3.88–11.88)	-57.43	1.09 (0.54–1.97)	-47.24	6.91 (3.65–11.12)	-52.96	1.09 (0.52–1.95)	-58.23
Syria	0.33 (0.22–0.49)	-75.16	0.28 (0.18–0.43)	-72.32	0.53 (0.36–0.75)	-72.64	0.13 (0.07–0.20)	-73.98
Tunisia	0.17 (0.10–0.27)	-77.38	0.12 (0.06–0.22)	-60.52	0.21 (0.13–0.32)	-72.55	0.07 (0.04–0.13)	-75.21
United Arab Emirates	0.08 (0.04–0.14)	-78.97	0.05 (0.02–0.09)	-73.71	0.04 (0.02–0.07)	-77.92	0.03 (0.01–0.05)	-75.71
Yemen	3.77 (1.79–7.49)	-60.35	1.15 (0.48–2.43)	-55.61	3.28 (1.64–6.14)	-61.43	0.90 (0.42–1.90)	-58.58

Table 3 Socio-demographic Index (SDI); years of life lost (YLL), years lived with disability (YLD), and disability-adjusted life years (DALY) rates and percent change; and YLL/YLD ratio for maternal causes in the Eastern Mediterranean Region, women ages 15–49, 1990–2015 (Global Burden of Disease 2015 study, Eastern Mediterranean countries, 1990–2015)

Country	SDI		YLL rate (per 100,000)		YLD rate (per 100,000)		YLL/YLD ratio		DALY rate (per 100,000)		% change
	1990	2015	% change	1990	2015	% change	1990	2015	1990	2015	
Eastern Mediterranean Region	0.55	600 (512–694)	-41	39 (27–52)	22 (15–30)	-43	15.38	16.09	639 (549–735)	376 (303–457)	-41
Afghanistan	0.29	2036 (1253–3135)	-30	26 (17–37)	16 (10–23)	-39	78.31	89.44	2062 (1274–3160)	1446 (856–2233)	-30
Bahrain	0.78	85 (63–116)	-78	14 (9–21)	6 (4–10)	-56	6.07	3.17	99 (77–132)	25 (19–33)	-75
Djibouti	0.46	869 (357–1586)	-22	68 (45–98)	48 (31–68)	-30	12.78	14.19	937 (424–1659)	729 (319–1941)	-22
Egypt	0.62	274 (229–319)	-76	15 (9–21)	11 (7–17)	-26	18.27	5.91	289 (245–333)	76 (61–94)	-74
Iran	0.72	104 (71–146)	-81	15 (9–22)	7 (4–10)	-56	6.93	2.71	119 (84–162)	26 (19–37)	-78
Iraq	0.58	312 (226–417)	-65	17 (10–24)	13 (8–20)	-21	18.35	8.38	329 (241–436)	122 (75–190)	-63
Jordan	0.7	196 (149–252)	-82	20 (13–30)	14 (8–22)	-31	9.80	2.50	216 (168–271)	49 (37–61)	-77
Kuwait	0.86	10 (9–13)	-43	6 (4–9)	5 (3–8)	-9	1.67	1.20	16 (14–20)	11 (9–15)	-30
Lebanon	0.75	51 (33–79)	-74	10 (6–15)	5 (3–8)	-47	5.10	2.60	61 (42–90)	18 (13–27)	-70
Libya	0.64	41 (29–56)	-42	12 (8–18)	8 (5–13)	-34	3.42	3.00	54 (41–69)	32 (23–43)	-40
Morocco	0.5	541 (418–714)	-86	15 (9–22)	9 (5–14)	-41	36.07	8.22	556 (432–730)	83 (54–127)	-85
Oman	0.73	71 (42–113)	-78	18 (11–26)	7 (4–12)	-59	3.94	2.14	88 (58–131)	23 (17–30)	-74
Pakistan	0.47	910 (696–1142)	-36	74 (50–102)	41 (28–57)	-45	12.30	14.17	984 (777–1213)	622 (469–784)	-37
Palestine	0.57	76 (50–113)	-62	19 (12–28)	12 (7–18)	-38	4.00	2.42	95 (66–132)	41 (32–55)	-57
Qatar	0.8	77 (51–110)	-79	10 (6–15)	4 (3–7)	-55	7.70	4.00	87 (60–120)	21 (14–27)	-76
Saudi Arabia	0.76	42 (32–55)	-61	16 (10–24)	8 (5–12)	-49	2.63	2.00	58 (46–72)	25 (20–30)	-58
Somalia	0.15	2144 (361–5693)	-19	102 (68–149)	61 (42–87)	-40	21.02	28.54	2246 (464–5797)	1802 (409–5273)	-20
Sudan	0.43	1159 (758–1561)	-55	43 (28–64)	26 (17–36)	-41	26.95	20.23	1202 (803–1602)	552 (306–810)	-54
Syria	0.58	264 (199–341)	-72	17 (11–26)	10 (6–16)	-42	15.53	7.40	281 (216–357)	84 (65–106)	-70
Tunisia	0.65	128 (100–163)	-69	11 (7–17)	7 (4–11)	-36	11.64	5.57	140 (110–175)	47 (33–63)	-67
United Arab Emirates	0.88	44 (28–68)	-77	11 (7–16)	3 (2–5)	-72	4.00	3.33	55 (37–79)	13 (9–19)	-76
Yemen	0.41	1235 (656–1865)	-55	104 (73–141)	42 (29–59)	-60	11.88	13.38	1339 (754–1982)	604 (347–1103)	-55

Fig. 2 Disability-adjusted life-year (DALY) rates per 100,000 women ages 15–49 for maternal causes by country, 2015 (Global Burden of Disease 2015 Study, Eastern Mediterranean Countries 2015)





Fig. 3 Disability-adjusted life-year (DALY) rates per 100,000 women ages 10–54 for maternal causes in the Eastern Mediterranean Region by age, 2015 (Global Burden of Disease 2015 study, Eastern Mediterranean Region 2015)



implemented to improve determinants of health and health coverage. Increases in the number of health facilities and training programs led to the percentage of deliveries assisted by skilled personnel increasing from 31% in 1992 to 73.5% in 2011. Other successes in the region occurred in Jordan and Iran, where maternal mortality was reduced by more than 50% from levels in 1990 (Kassebaum et al. 2016b).

Uneven improvements in MMR in the region may be related to differing levels of ramp-up in coverage for specific types of reproductive health care—antenatal care, in-facility delivery, skilled birth attendance, family

planning services, emergency obstetric care, and post-natal care—that are all known to decrease the risk of negative pregnancy outcomes (Lim et al. 2010; Randive et al. 2014). Quality of care must also be considered during buildup of the maternal health care system. Programs for reproductive health care must ensure that women are receiving the care they need during pregnancy and the post-partum period (Rowe et al. 2005). Care should be integrated and not be focused on single vertical interventions (Campbell et al. 2006).

It has been challenging for many countries to produce timely and accurate data on levels of maternal mortality

that would indicate the extent of their progress in reducing maternal deaths (Mokdad et al. 2016). Reliable information is a necessary component of any strategy aimed at reducing maternal mortality. Continued progress in data collection in the EMR is key to evaluating progress in reducing maternal mortality.

Our study has several limitations. First, many countries in the region have poor health data and vital statistics. We used GBD methodology to account for quality and lack of data. We also applied our standard GBD garbage code correction to address this limitation. Second, little information is available on unsafe abortion in the region due to religion and culture. However, our study is the most comprehensive on burden of diseases and applies the standard methodology that allows global comparison.

Conclusion

Progress in reducing maternal mortality in the EMR has accelerated in the past 15 years, but there is still much to do to reduce preventable deaths. Extending basic maternal health services, improving quality of care, and eliminating unmet need for contraception are all crucial, proven steps effective at reducing MMR. Our study showed the importance of empowering women: increased women's rights are needed to improve their health. Finally, coordinated and rigorous efforts are needed to make sure that every woman in need receives these interventions in a timely fashion at each stage of her reproductive life.

GBD 2015 Eastern Mediterranean Region Maternal Mortality

Collaborators: Ibrahim Khalil, MD, Institute for Health Metrics and Evaluation, University of Washington. Michael Collison, BS, Institute for Health Metrics and Evaluation, University of Washington, Seattle, Washington, United States. Charbel El Bcheraoui, PhD, Institute for Health Metrics and Evaluation, University of Washington. Raghid Charara, MD, American University of Beirut, Beirut, Lebanon. Maziar Moradi-Lakeh, MD, Department of Community Medicine, Preventative Medicine and Public Health Research Center, Gastrointestinal and Liver Disease Research Center (GILDRC), University of Medical Sciences, Tehran, Iran. Ashkan Afshin MD, Institute for Health Metrics and Evaluation, University of Washington, Seattle, Washington, United States. Adrienne Chew, ND, Institute for Health Metrics and Evaluation, University of Washington. Farah Daoud, BA/BS, Institute for Health Metrics and Evaluation, University of Washington. Kristopher J. Krohn, BA, Institute for Health Metrics and Evaluation, University of Washington, Seattle, Washington, United States. Danny Colombara, PhD, Institute for Health Metrics and Evaluation, University of Washington, Seattle, Washington, United States. Rebecca Ehrenkranz, MPH, Institute for Health Metrics and Evaluation, University of Washington, Seattle, Washington, United States. Michael Kutz, BS, Institute for Health Metrics and Evaluation, University of Washington, Seattle, Washington, United States. Haidong Wang, PhD, Institute for Health Metrics and Evaluation, University of Washington, Seattle, Washington, United States. Amanuel Alemu Abajobir, MPH, School of Public Health, University of Queensland, Brisbane, QLD, Australia. Foad Abd-Allah, MD,

Department of Neurology, Cairo University, Cairo, Egypt. Haftom Niguse Abraha, MS, Mekelle University, Mekelle, Ethiopia. Laith J. Abu-Raddad, PhD, Infectious Disease Epidemiology Group, Weill Cornell Medical College in Qatar, Doha, Qatar. Aliasghar Ahmad Kiadaliri, PhD, Department of Clinical Sciences Lund, Orthopedics, Clinical Epidemiology Unit, Lund University, Lund, Sweden. Alireza Ahmadi, PhD, Kermanshah University of Medical Sciences, Kermanshah, Iran. Kedir Yimam Ahmed, MPH, Debre Markos University, Debre Markos, Ethiopia. Muktar Beshir Ahmed, MPH, College of Health Sciences, Department of Epidemiology, ICT and e-Learning Coordinator, Jimma University, Jimma, Ethiopia. Faris Hasan Al Lami, PhD, Baghdad College of Medicine, Baghdad, Iraq. Khurshid Alam, PhD, Murdoch Childrens Research Institute, The University of Melbourne, Parkville, Victoria, Australia; The University of Melbourne, Melbourne, VIC, Australia; The University of Sydney, Sydney, NSW, Australia. Deena Alasfoor, MSc, Ministry of Health, Al Khuwair, Oman. Reza Alizadeh-Navaei, PhD, Gastrointestinal Cancer Research Center, Mazandaran University of Medical Sciences, Sari, Iran. Juma M Alkaabi, MD, College of Medicine and Health Sciences United Arab Emirates University, Al-Ain City, United Arab Emirates. Fatma Al-Maskari, PhD, College of Medicine & Health Sciences, United Arab Emirates; University, Al-Ain City, United Arab Emirates. Rajaa Al-Raddadi, PhD, Joint Program of Family and Community Medicine, Jeddah, Saudi Arabia. Khalid A. Altirkawi, MD, King Saud University, Riyadh, Saudi Arabia. Nahla Anber, PhD, Mansoura University, Mansoura, Egypt. Hossein Ansari, PhD, Health Promotion Research Center, Department of Epidemiology and Biostatistics, Zahedan University of Medical Sciences, Zahedan, Iran. Hamid Asayesh, PhD, Department of Medical Emergency, School of Paramedic, Qom University of Medical Sciences, Qom, Iran. Rana Jawad Asghar, MD, South Asian Public Health Forum, Islamabad, Pakistan. Tesfay Mehari Atey, MS, Mekelle University, Mekelle, Ethiopia. Tadesse Awoke Ayele, MS, University of Gondar, Gondar, Ethiopia. Till Bärnighausen, MD, Department of Global Health and Population, Harvard T. H. Chan School of Public Health, Harvard University, Boston, MA, United States; Africa Health Research Institute, Mtubatuba, KwaZulu-Natal, South Africa; Institute of Public Health, Heidelberg University, Heidelberg, Germany. Umar Bacha, PhD, School of Health Sciences, University of Management and Technology, Lahore, Pakistan. Aleksandra Barac, PhD, Faculty of Medicine, University of Belgrade, Belgrade, Serbia. Suzanne L. Barker-Collo, PhD, School of Psychology, University of Auckland, Auckland, New Zealand. Bernhard T. Baune, PhD, School of Medicine, University of Adelaide, Adelaide, South Australia, Australia. Shahrzad Bazargan-Hejazi, PhD, College of Medicine, Charles R. Drew University of Medicine and Science, Los Angeles, CA, United States; David Geffen School of Medicine, University of California at Los Angeles, Los Angeles, CA, United States. Neeraj Bedi, MD, College of Public Health and Tropical Medicine, Jazan, Saudi Arabia. Isabela M Bensenor, PhD, University of São Paulo, São Paulo, Brazil. Adugnaw Berhane, PhD, College of Health Sciences, Debre Berhan University, Debre Berhan, Ethiopia. Addisu Shunu Beyene, MPH, College of Health and Medical Science, Haramaya University, Harar, Ethiopia. Zulfiqar A. Bhutta, PhD, Centre of Excellence in Women and Child Health, Aga Khan University, Karachi, Pakistan; Centre for Global Child Health, The Hospital for Sick Children, Toronto, ON, Canada. Dube Jara Boneya, MPH, Department of Public Health, Debre Markos University, Debre Markos, Ethiopia. Rohan Borschmann, PhD, The University of Melbourne, Melbourne, VIC, Australia; Murdoch Childrens Research Institute, Melbourne, VIC, Australia. Nicholas J. K. Breitborde, PhD, The Ohio State University, Columbus, OH, United States. Zahid A. Butt, PhD, Al Shifa Trust Eye Hospital, Rawalpindi, Punjab, Pakistan. Ferrán Catalá-López, PhD, Department of Medicine, University of Valencia/INCLIVA Health Research Institute and CIBERSAM, Valencia, Spain; Clinical Epidemiology Program, Ottawa Hospital Research Institute, Ottawa,

Canada. Liliana G. Ciobanu, MS, School of Medicine, University of Adelaide, Adelaide, South Australia, Australia. Hadi Danawi, PhD, Walden University, Minneapolis, Minnesota, United States. Amare Deribew, PhD, Nuffield Department of Medicine, University of Oxford, Oxford, United Kingdom; KEMRI-Wellcome Trust Research Programme, Kilifi, Kenya. Samath D. Dharmaratne, MD, Department of Community Medicine, Faculty of Medicine, University of Peradeniya, Peradeniya, Sri Lanka. Kerrie E. Doyle, PhD, RMIT University, Bundoora, VIC, Australia; Australian National University, Canberra, ACT, Australia. Aman Yesuf Endries, MPH, Arba Minch University, Arba Minch, Ethiopia. Emerito Jose Aquino Faraon, MD, College of Public Health, University of the Philippines Manila, Manila, Philippines; Department of Health, Manila, Philippines. André Faro, PhD, Federal University of Sergipe, Aracaju, Brazil. Maryam S. Farvid, PhD, Department of Nutrition, Harvard T. H. Chan School of Public Health, Harvard University, Boston, MA, United States; Harvard/MGH Center on Genomics, Vulnerable Populations, and Health Disparities, Mongan Institute for Health Policy, Massachusetts General Hospital, Boston, MA, United States. Wubalem Fekadu, MS, Bahir Dar University, Bahir Dar, Ethiopia. Seyed-Mohammad Fereshtehnejad, PhD, Department of Neurobiology, Care Sciences and Society (NVS), Karolinska Institutet, Stockholm, Sweden. Florian Fischer, PhD, School of Public Health, Bielefeld University, Bielefeld, Germany. Tsegaye Tewelde Gebrehiwot, MPH, Jimma University, Jimma, Ethiopia. Ababi Zergaw Giref, PhD, Addis Ababa University, Addis Ababa, Ethiopia. Melkamu Dedefo Gishu, MS, Haramaya University, Dire Dawa, Ethiopia; Kersa Health and Demographic Surveillance System, Harar, Ethiopia. Alessandra Carvalho Goulart, PhD, Center for Clinical and Epidemiological Research Center- Hospital Universitario-University of São Paulo, São Paulo, Brazil; Center of Check of Hospital Sirio Libanes, São Paulo, Brazil. Tesfa Dejenie Habtewold, MS, University of Groningen, Groningen, Netherlands; Debre Berhan University, Debre Berhan, Ethiopia. Randah Ribhi Hamadeh, DPhil, Arabian Gulf University, Manama, Bahrain. Mitiku Teshome Hambisa, MPH, College of Health and Medical Sciences, Haramaya University, Harar, Ethiopia. Samer Hamidi, DrPH, Hamdan Bin Mohammed Smart University, Dubai, United Arab Emirates. Josep Maria Haro, MD, Parc Sanitari Sant Joan de Déu - CIBERSAM, Sant Boi de Llobregat (Barcelona), Spain. Mohammad Sadegh Hassanvand, PhD, Center for Air Pollution Research, Institute for Environmental Research, Tehran University of Medical Sciences, Tehran, Iran. Nobuyuki Horita, MD, Department of Pulmonology, Yokohama City University Graduate School of Medicine, Yokohama, Kanagawa, Japan. Mohamed Hsairi, MD, Department of Epidemiology, Salah Azaiz Institute, Tunis, Tunisia. Hsiang Huang, MD, Cambridge Health Alliance, Cambridge, MA, United States. Abdullatif Husseini, PhD, Institute of Community and Public Health, Birzeit University, Birzeit, Palestine. Mihajlo B. Jakovljevic, PhD, Faculty of Medical Sciences, University of Kragujevac, Kragujevac, Central Serbia - Sumadija, Serbia; The Center for Health Trends and Forecasts, Institute for Health Metrics and Evaluation, University of Washington, Seattle, Washington, United States. Spencer Lewis James, MD, Denver Health/University of Colorado, Denver, CO, United States. Jost B. Jonas, MD, Department of Ophthalmology, Medical Faculty Mannheim, Ruprecht-Karls-University Heidelberg, Mannheim, Germany. Amir Kasaian, PhD, Hematology-Oncology and Stem Cell Transplantation Research Center, Tehran University of Medical Sciences, Tehran, Iran; Endocrinology and Metabolism Population Sciences Institute, Tehran University of Medical Sciences, Tehran, Iran. Yousef Saleh Khader, ScD, Department of Community Medicine, Public Health and Family Medicine, Jordan University of Science and Technology, Irbid, Jordan. Ejaz Ahmad Khan, MD, Health Services Academy, Islamabad, Pakistan. Abdullah Tawfih Abdullah Khoja, MD, Mohammed Ibn Saudi University, Riyadh, Saudi Arabia. Ardashir Khosravi, PhD, Iranian Ministry of Health and Medical Education, Tehran, Iran, Non-communicable Diseases Research Center, Tehran University of Medical Sciences, Tehran, Iran. Jagdish Khubchandani, PhD, Department of Nutrition and Health Science, Ball State University, Muncie, Indiana, United States. Daniel Kim, DrPH, Department of Health Sciences, Northeastern University, Boston, Massachusetts, United States. Yun Jin Kim, PhD, Faculty of Chinese Medicine, Southern University College, Skudai, Malaysia. Yoshihiro Kokubo, PhD, Department of Preventive Cardiology, National Cerebral and Cardiovascular Center, Suita, Japan. Ai Koyanagi, MD, Research and Development Unit, Parc Sanitari Sant Joan de Deu (CIBERSAM), Barcelona, Spain. Barthelemy Kuate Defo, PhD, Department of Social and Preventive Medicine, School of Public Health, University of Montreal, Montreal, Quebec, Canada; Department of Demography and Public Health Research Institute, University of Montreal, Montreal, Canada. Heidi J. Larson, PhD, Department of Infectious Disease Epidemiology, London School of Hygiene & Tropical Medicine, London, United Kingdom; Institute for Health Metrics and Evaluation, University of Washington, Seattle, Washington, United States. Asma Abdul Latif, PhD, Department of Zoology, Lahore College for Women University, Lahore, Punjab, Pakistan. Paul H Lee, PhD, Hong Kong Polytechnic University, Hong Kong, China. Cheru Tesema Leshargie, MPH, Debre Markos University, Debre Markos, Ethiopia. Ricky Leung, PhD, State University of New York, Albany, Rensselaer, NY, United States. Loon-Tzian Lo, MD, UnionHealth Associates, LLC, St. Louis, MO, United States; Alton Mental Health Center, Alton, IL, United States. Raimundas Lunevicius, PhD, Aintree University Hospital National Health Service Foundation Trust, Liverpool, United Kingdom; School of Medicine, University of Liverpool, Liverpool, United Kingdom. Hassan Magdy Abd El Razek, MBChB, Mansoura Faculty of Medicine, Mansoura, Egypt. Mohammed Magdy Abd El Razek, MBChB, Aswan University Hospital, Aswan Faculty of Medicine, Aswan, Egypt. Reza Majdzadeh, PhD, Knowledge Utilization Research Center and Community Based Participatory Research Center, Tehran University of Medical Sciences, Tehran, Iran. Azeem Majeed, MD, Department of Primary Care & Public Health, Imperial College London, London, United Kingdom. Reza Malekzadeh, MD, Digestive Diseases Research Institute, Tehran University of Medical Sciences, Tehran, Iran; Digestive Diseases Research Institute, Tehran University of Medical Sciences, Tehran, Iran. Jose Martinez-Raga, PhD, Hospital Universitario Doctor Peset, Valencia, Spain; CEU Cardinal Herrera University, Moncada, Spain. Habibolah Masoudi Farid, MD, State Welfare Organisation, Tehran, Iran. Mohsen Mazidi, PhD, Key State Laboratory of Molecular Developmental Biology, Institute of Genetics and Developmental Biology, Chinese Academy of Sciences, Beijing, China. John J. McGrath, PhD, Queensland Centre for Mental Health Research, The Park Centre for Mental Health, Wacol, Queensland, Australia; Queensland Brain Institute, University of Queensland, St. Lucia, QLD, Australia, National Centre for Register-Based Research, Aarhus School of Business and Social Sciences, Aarhus University, Aarhus, Denmark. Ziad A. Memish, MD, Saudi Ministry of Health, Riyadh, Saudi Arabia; College of Medicine, Alfaisal University, Riyadh, Saudi Arabia. Walter Mendoza, MD, United Nations Population Fund, Lima, Peru. Melkamu Merid Mengesha, MPH, College of Health and Medical Sciences, Haramaya University, Harar, Ethiopia. Mubarek Abera Mengistie, MS, Jimma University, Jimma, Ethiopia. George A. Mensah, MD, Center for Translation Research and Implementation Science, National Heart, Lung, and Blood Institute, National Institutes of Health, Bethesda, MD, United States. Haftay Berhane Mezgebe, MS, Mekelle University, Mekelle, Ethiopia. Ted R. Miller, PhD, Pacific Institute for Research & Evaluation, Calverton, MD, United States, Centre for Population Health, Curtin University, Perth, WA, Australia. Philip B. Mitchell, MD, University of New South Wales, Kensington, New South Wales, Australia. Alireza Mohammadi, PhD, Neuroscience Research Center, Baqiyatallah University of Medical Science,

- Tehran, Iran. Shafiu Mohammed, PhD, Health Systems and Policy Research Unit, Ahmadu Bello University, Zaria, Nigeria; Institute of Public Health, Heidelberg University, Heidelberg, Germany. Carla Makhoul Obermeyer, DSc, Center for Research on Population and Health, Faculty of Health Sciences, American University of Beirut, Beirut, Lebanon. Felix Akpojene Ogbo, MPH, Centre for Health Research, Western Sydney University, Sydney, New South Wales, Australia. Elizabeth Palomares Castillo, PhD, Ministry of Health, Mexico City, Mexico City, Mexico; Universidad Nacional Autónoma de México (UNAM), Mexico City, Mexico City, Mexico. Christina Papachristou, PhD, Charité University Medicine Berlin, Berlin, Germany. Scott B. Patten, PhD, Department of Community Health Sciences, University of Calgary, Calgary, Alberta, Canada. George C. Patton, MD, Murdoch Childrens Research Institute, Department of Paediatrics, University of Melbourne, Melbourne, Victoria, Australia. David M Pereira, PhD, REQUIMTE/LAQV, Laboratório de Farmacognosia, Departamento de Química, Faculdade de Farmácia, Universidade do Porto, Porto, Portugal. Aslam Pervaiz, MHA, Postgraduate Medical Institute, Lahore, Pakistan. Michael Robert Phillips, MD, Shanghai Jiao Tong University School of Medicine, Shanghai, China; Emory University, Atlanta, Georgia, United States. Farshad Pourmalek, PhD, University of British Columbia, Vancouver, British Columbia, Canada. Mostafa Qorbani, PhD, Non-communicable Diseases Research Center, Alborz University of Medical Sciences, Karaj, Iran. Amir Radfar, MD, A T Still University, Kirksville, MO, United States. Anwar Rafay, MS, Contech International Health Consultants, Lahore, Pakistan, Contech School of Public Health, Lahore, Pakistan. Vafa Rahimi-Movaghar, MD, Sina Trauma and Surgery Research Center, Tehran University of Medical Sciences, Tehran, Iran. Rajesh Kumar Rai, MPH, Society for Health and Demographic Surveillance, Suri, India. David Laith Rawaf, MD, WHO Collaborating Centre, Imperial College London, London, United Kingdom; North Hampshire Hospitals, Basingstoke, United Kingdom; University College London Hospitals, London, United Kingdom. Salman Rawaf, MD, Imperial College London, London, United Kingdom. Amany H. Refaat, PhD, Walden University, Minneapolis, MN, United States; Suez Canal University, Ismailia, Egypt. Satar Rezaei, PhD, School of Public Health, Kermanshah University of Medical Sciences, Kermanshah, Iran. Mohammad Sadeq Rezai, MD, Mazandaran University of Medical Sciences, Sari, Iran. Gholamreza Roshandel, PhD, Golestan Research Center of Gastroenterology and Hepatology, Golestan University of Medical Sciences, Gorgan, Iran; Digestive Diseases Research Institute, Tehran University of Medical Sciences, Tehran, Iran. Mahdi Safdarian, MD, Sina Trauma & Surgery Research Center, Tehran University of Medical Sciences, Tehran, Iran. Mahdi Safiabadi, MD, Student Research Committee, Baqiyatallah University of Medical Sciences, Tehran, Iran. Saeid Safiri, PhD, Managerial Epidemiology Research Center, Department of Public Health, School of Nursing and Midwifery, Maragheh University of Medical Sciences, Maragheh, Iran. Rajesh Sagar, MD, All India Institute of Medical Sciences, New Delhi, India. Mohammad Ali Sahraian, MD, MS Research Center, Neuroscience Institute, Tehran University of Medical Sciences, Tehran, Iran. Payman Salamati, MD, Sina Trauma and Surgery Research Center, Tehran University of Medical Sciences, Tehran, Iran. Abdallah M. Samy, PhD, Ain Shams University, Cairo, Egypt, Lawrence, Kansas, United States. Benn Sartorius, PhD, Public Health Medicine, School of Nursing and Public Health, University of KwaZulu-Natal, Durban, South Africa; UKZN Gastrointestinal Cancer Research Centre, South African Medical Research Council (SAMRC), Durban, South Africa. Mete I. Saylan, PhD, Bayer Turkey, Istanbul, Turkey. Soraya Seedat, PhD, Stellenbosch University, Cape Town, South Africa. Sadaf G. Sepanlou, PhD, Digestive Diseases Research Institute, Tehran University of Medical Sciences, Tehran, Iran. Masood Ali Shaikh, MD, Independent Consultant, Karachi, Pakistan. Morteza Shamzadeh, MPH, Department of Medical Surgical Nursing, School of Nursing and Midwifery, Hamadan University of Medical Sciences, Hamadan, Iran. Diego Augusto Santos Silva, PhD, Federal University of Santa Catarina, Florianopolis, Brazil. Jasvinder A. Singh, MD, University of Alabama at Birmingham and Birmingham Veterans Affairs Medical Center, Birmingham, Alabama, United States. Badr H. A. Sobaih, MD, King Saud University, Riyadh, Saudi Arabia. Chandrashekhar T. Sreeramareddy, MD, Department of Community Medicine, International Medical University, Kuala Lumpur, Malaysia. Dan J. Stein, PhD, Department of Psychiatry, University of Cape Town, Cape Town, South Africa; South African Medical Research Council Unit on Anxiety & Stress Disorders, Cape Town, South Africa. Rizwan Suliankatchi Abdulkader, MD, Ministry of Health, Kingdom of Saudi Arabia, Riyadh, Saudi Arabia. Bryan L. Sykes, PhD, Departments of Criminology, Law & Society, Sociology, and Public Health, University of California, Irvine, Irvine, CA, United States. Rafael Tabarés-Seisdedos, PhD, Department of Medicine, University of Valencia, INCLIVA Health Research Institute and CIBERSAM, Valencia, Spain. Muawiyah Sufiyan, MD, Ahmadu Bello University, Zaria, Nigeria. Karen M. Tabb, PhD, School of Social Work, University of Illinois at Urbana-Champaign, Champaign, Illinois, United States. Arash Tehrani-Banihashemi, PhD, Preventive Medicine and Public Health Research Center, Iran University of Medical Sciences, Tehran, Iran. Mohamad-Hani Temsah, MD, King Saud University, Riyadh, Saudi Arabia. Abdullah Sulieman Terkawi, MD, Department of Anesthesiology, University of Virginia, Charlottesville, VA, United States; Department of Anesthesiology, King Fahad Medical City, Riyadh, Saudi Arabia. Outcomes Research Consortium, Cleveland Clinic, Cleveland, OH, United States. Roman Topor-Madry, PhD, Institute of Public Health, Faculty of Health Sciences, Jagiellonian University Medical College, Kraków, Poland; Faculty of Health Sciences, Wrocław Medical University, Wrocław, Poland. Kingsley Nnana Ukwaja, MD, Department of Internal Medicine, Federal Teaching Hospital, Abakaliki, Nigeria. Olalekan A. Uthman, PhD, Warwick Medical School, University of Warwick, Coventry, United Kingdom. Stein Emil Vollset, DrPH, Center for Disease Burden, Norwegian Institute of Public Health, Bergen, Norway; Department of Global Public Health and Primary Care, University of Bergen, Bergen, Norway, Institute for Health Metrics and Evaluation, University of Washington, Seattle, Washington, United States. Tolassa Wakayo, MS, Jimma University, Jimma, Ethiopia. Yuan-Pang Wang, PhD, University of São Paulo Medical School, São Paulo, Brazil. Andrea Werdecker, PhD, Competence Center Mortality-Follow-Up of the German National Cohort, Federal Institute for Population Research, Wiesbaden, Germany. Ronny Westerman, PhD, Federal Institute for Population Research, Wiesbaden, Germany; German National Cohort Consortium, Heidelberg, Germany. Abdulhalik Workicho, MPH, Jimma University, Jimma, Ethiopia; Ghent University, Ghent, Belgium. Mohsen Yaghoubi, MSc, School of Public Health, University of Saskatchewan, Saskatoon, Saskatchewan, Canada. Hassen Hamid Yimam, MPH, Mizan Tepi University, Mizan Teferi, Ethiopia. Naohiro Yonemoto, MPH, Department of Biostatistics, School of Public Health, Kyoto University, Kyoto, Japan. Mustafa Z Younis, DrPH, Jackson State University, Jackson, MS, United States. Chuanhua Yu, PhD, Department of Epidemiology and Biostatistics, School of Public Health, Wuhan University, Wuhan, China, Global Health Institute, Wuhan University, Wuhan, China. Maysaa El Sayed Zaki, PhD, Faculty of Medicine, Mansoura University, Mansoura, Egypt. Aisha O. Jumaan, PhD, Independent Consultant, Seattle, Washington, United States. Theo Vos, PhD, Institute for Health Metrics and Evaluation, University of Washington, Seattle, Washington, United States. Simon I. Hay, DSc, Oxford Big Data Institute, Li Ka Shing Centre for Health Information and Discovery, University of Oxford, Oxford, United Kingdom; Institute for Health Metrics and Evaluation, University of Washington, Seattle, Washington, United States. Mohsen Naghavi, PhD, Institute for Health Metrics and

Evaluation, University of Washington, Seattle, Washington, United States. Nicholas J. Kassebaum, MD, Institute for Health Metrics and Evaluation, University of Washington, Seattle, Washington, United States; Department of Anesthesiology & Pain Medicine, Seattle Children's Hospital, Seattle, Washington, United States. Christopher J. L. Murray, DPhil, Institute for Health Metrics and Evaluation, University of Washington, Seattle, Washington, United States. Ali H. Mokdad, PhD, Institute for Health Metrics and Evaluation, University of Washington, Seattle, Washington, United States.

Compliance with ethical standards

This manuscript reflects original work that has not previously been published in whole or in part and is not under consideration elsewhere. All authors have read the manuscript and have agreed that the work is ready for submission and accept responsibility for its contents. The authors of this paper have complied with all ethical standards and do not have any conflicts of interest to disclose at the time of submission. The funding source played no role in the design of the study, the analysis and interpretation of data, and the writing of the paper. The study did not involve human participants and/or animals; therefore, no informed consent was needed.

Funding This research was funded by the Bill & Melinda Gates Foundation.

Conflict of interest The authors declare that they have no conflicts of interest at this time.

Open Access This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

References

- Abouzahir C, Wardlaw T (2001) Maternal mortality at the end of a decade: signs of progress? *Bull World Health Org* 79:561–573
- Alvarez JL, Gil R, Hernández V, Gil A (2009) Factors associated with maternal mortality in Sub-Saharan Africa: an ecological study. *BMC Public Health* 9:462. doi:10.1186/1471-2458-9-462
- Bayati M, Akbarian R, Kavosi Z (2013) Determinants of life expectancy in eastern mediterranean region: a health production function. *Int J Health Policy Manag* 1:57–61. doi:10.15171/ijhpm.2013.09
- Buor D, Bream K (2004) An analysis of the determinants of maternal mortality in sub-Saharan Africa. *J Womens Health* 13:926–938. doi:10.1089/jwh.2004.13.926
- Campbell OMR, Graham WJ, Lancet Maternal Survival Series steering group (2006) Strategies for reducing maternal mortality: getting on with what works. *Lancet Lond Engl* 368:1284–1299. doi:10.1016/S0140-6736(06)69381-1
- Hertz E, Hebert JR, Landon J (1994) Social and environmental factors and life expectancy, infant mortality, and maternal mortality rates: results of a cross-national comparison. *Soc Sci Med* 39:105–114. doi:10.1016/0277-9536(94)90170-8
- Homaie Rad E, Vahedi S, Teimourizad A et al (2013) Comparison of the effects of public and private health expenditures on the health status: a panel data analysis in eastern mediterranean countries. *Int J Health Policy Manag* 1:163–167. doi:10.15171/ijhpm.2013.29
- Institute for Health Metrics and Evaluation (2016) Global health data exchange | GHDx. <http://ghdx.healthdata.org/>. Accessed 1 June 2017
- Institute for Health Metrics and Evaluation (2017) Global burden of disease data visualization. <https://vizhub.healthdata.org/gbd-compare/>. Accessed 13 July 2017
- Kassebaum NJ, Bertozzi-Villa A, Coggeshall MS et al (2014) Global, regional, and national levels and causes of maternal mortality during 1990–2013: a systematic analysis for the Global Burden of Disease Study 2013. *Lancet* 384:980–1004. doi:10.1016/S0140-6736(14)60696-6
- Kassebaum NJ, Arora M, Barber RM et al (2016a) Global, regional, and national disability-adjusted life-years (DALYs) for 315 diseases and injuries and healthy life expectancy (HALE), 1990–2015: a systematic analysis for the Global Burden of Disease Study 2015. *Lancet* 388:1603–1658. doi:10.1016/S0140-6736(16)31460-X
- Kassebaum NJ, Barber RM, Bhutta ZA et al (2016b) Global, regional, and national levels of maternal mortality, 1990–2015: a systematic analysis for the Global Burden of Disease Study 2015. *Lancet* 388:1775–1812. doi:10.1016/S0140-6736(16)31470-2
- Liang J, Zhu J, Dai L et al (2010) Maternal mortality in China, 1996–2005. *Int J Gynecol Obstet* 110:93–96. doi:10.1016/j.ijgo.2010.03.013
- Lim SS, Dandona L, Hoisington JA et al (2010) India's Janani Suraksha Yojana, a conditional cash transfer programme to increase births in health facilities: an impact evaluation. *Lancet* 375:2009–2023. doi:10.1016/S0140-6736(10)60744-1
- Midhet F, Becker S, Berendes HW (1998) Contextual determinants of maternal mortality in rural Pakistan. *Soc Sci Med* 46:1587–1598
- Mokdad AH, Forouzanfar MH, Daoud F et al (2016) Health in times of uncertainty in the eastern Mediterranean region, 1990–2013: a systematic analysis for the Global Burden of Disease Study 2013. *Lancet Glob Health* 4:e704–713. doi:10.1016/S2214-109X(16)30168-1
- Muldoon KA, Galway LP, Nakajima M et al (2011) Health system determinants of infant, child and maternal mortality: a cross-sectional study of UN member countries. *Glob Health* 7:42. doi:10.1186/1744-8603-7-42
- Peterson HB, Haidar J, Merialdi M et al (2012) Preventing maternal and newborn deaths globally: using innovation and science to address challenges in implementing life-saving interventions. *Obstet Gynecol* 120:636–642. doi:10.1097/AOG.0b013e3182632cc1
- Randive B, San Sebastian M, De Costa A, Lindholm L (2014) Inequalities in institutional delivery uptake and maternal mortality reduction in the context of cash incentive program, Janani Suraksha Yojana: results from nine states in India. *Soc Sci Med* 123:1–6. doi:10.1016/j.socscimed.2014.10.042
- Ribeiro PS, Jacobsen KH, Mathers CD, Garcia-Moreno C (2008) Priorities for women's health from the Global Burden of Disease study. *Int J Gynaecol Obstet Off Organ Int Fed Gynaecol Obstet* 102:82–90. doi:10.1016/j.ijgo.2008.01.025
- Ronsmans C, Graham WJ (2006) Maternal mortality: who, when, where, and why. *Lancet* 368:1189–1200. doi:10.1016/S0140-6736(06)69380-X
- Rowe AK, de Savigny D, Lanata CF, Victora CG (2005) How can we achieve and maintain high-quality performance of health workers in low-resource settings? *Lancet Lond Engl* 366:1026–1035. doi:10.1016/S0140-6736(05)67028-6
- Thaddeus S, Maine D (1994) Too far to walk: maternal mortality in context. *Soc Sci Med* 38:1091–1110
- UNICEF (2016) Female genital mutilation/cutting: a global concern. https://www.unicef.org/media/files/FGMC_2016_brochure_final_UNICEF_SPREAD.pdf. Accessed 13 July 2017

- United Nations (2015) Transforming our world: the 2030 agenda for sustainable development. <https://sustainabledevelopment.un.org/post2015/transformingourworld>. Accessed 12 July 2017
- Vos T, Allen C, Arora M et al (2016) Global, regional, and national incidence, prevalence, and years lived with disability for 310 diseases and injuries, 1990–2015: a systematic analysis for the Global Burden of Disease Study 2015. *Lancet* 388:1545–1602. doi:[10.1016/S0140-6736\(16\)31678-6](https://doi.org/10.1016/S0140-6736(16)31678-6)
- Wang H, Naghavi M, Allen C et al (2016) Global, regional, and national life expectancy, all-cause mortality, and cause-specific mortality for 249 causes of death, 1980–2015: a systematic analysis for the Global Burden of Disease Study 2015. *Lancet* 388:1459–1544. doi:[10.1016/S0140-6736\(16\)31012-1](https://doi.org/10.1016/S0140-6736(16)31012-1)
- Warren E, Post N, Hossain M et al (2015) Systematic review of the evidence on the effectiveness of sexual and reproductive health interventions in humanitarian crises. *BMJ Open*. doi:[10.1136/bmjopen-2015-008226](https://doi.org/10.1136/bmjopen-2015-008226)
- WHO (2017) High-quality healthcare needed for girls and women who have experienced female genital mutilation. <http://www.who.int/reproductivehealth/topics/fgm/en/>
- WHO, UNICEF, UNFPA, World Bank (2007) Maternal mortality in 2005. http://www.who.int/whosis/mme_2005.pdf. Accessed 13 July 2017
- World Health Organization (1992) International Statistical classification of diseases and related health problems. Tenth Revision

Electronic Supplementary Material

Article title:

Maternal mortality and morbidity burden in the Eastern Mediterranean Region: Findings from the Global Burden of Disease 2015 study

Journal:

International Journal of Public Health

Authors:

GBD 2015 Eastern Mediterranean Region Maternal Mortality Collaborators


Corresponding author:

Ali H. Mokdad

Institute for Health Metrics and Evaluation, University of Washington, Seattle, WA, United States

Email: mokdaa@uw.edu

e-Figure 1 - Percent of total maternal disability-adjusted life-years(DALYs) attributable to maternal causes in the Eastern Mediterranean by age, 2015. (Global Burden of Disease 2015 Study, Eastern Mediterranean Countries, 2015).



Title: Maternal mortality and morbidity burden in the Eastern Mediterranean Region: findings from the Global Burden of Disease 2015 study

Group name: GBD 2015 Eastern Mediterranean Region Maternal Mortality Collaborators:

Prefix	First Name:	Middle Name	Last Name:	Email Addr	Highest de	Department	Institution/Organization	Institution, Institution/Organiz	Institution/Organizat	Department 2	Institution/Organization 2	Institution, Institution, Institution, Department	Institution, Institution, Institution, Institution					
Dr.	Ibrahim		Khalil	ikhali@uw MD		Institute for Health Metrics and Evaluation	University of Washington	Seattle	Washington	United_States								
Mr.	Michael		Collison	mc314@u BS		Institute for Health Metrics and Evaluation	University of Washington	Seattle	Washington	United_States								
Dr.	Charbel		El Bcheraoui	charbel@u PhD		Institute for Health Metrics and Evaluation	University of Washington											
Dr.	Raghid		Charara	raghidc@ MD			American University of Beirut	Beirut		Lebanon								
						Department of Community Medicine, Preventative Medicine and Public Health Research Center, Gastrointestinal and Liver Disease Research Center (GILDRC),												
Dr.	Maziar		Moradi-Lakeh	mazmoradi MD			Iran University of Medical Sciences	Tehran		Iran								
Dr.	Ashkan		Afshin	ashkan.afst MD		Institute for Health Metrics and Evaluation	University of Washington	Seattle	Washington	United_States								
Ms.	Adrienne		Chew	apchew@u ND			University of Washington											
Ms.	Farah		Daoud	farahd@uv BA/BS		Institute for Health Metrics and Evaluation	University of Washington											
Mr.	Kristopher	J.	Krohn	kkrohn1@ U BA			University of Washington	Seattle	Washington	United_States								
Dr.	Fiona	J	Charlson	fiona_charl PhD		School of Public Health	University of Queensland	Brisbane	Queensland	Australia	Institute for Health Metrics and Evaluation	University of Washington	Seattle	United_States	Queensland Brisbane	Queensland Australia		
Prof.	Danny		Colombara	dvc2@uw.r PhD		Institute for Health Metrics and Evaluation	University of Washington	Seattle	Washington	United_States								
Ms.	Louisa		Degenhardt	l.degenhardt PhD		National Drug and Alcohol Research Centre	University of New South Wales	Sydney		Australia								
Ms.	Rebecca		Ehrenkranz	rehrenk1@ MPH		Institute for Health Metrics and Evaluation	University of Washington	Seattle	Washington	United_States								
Dr.	Holly	E	Erskine	holly_erskii PhD			Queensland Centre for Mental Health Research	Brisbane	QLD	Australia	School of Public Health	University of Queensland	Brisbane	QLD	Australia	Institute fo University i Seattle	WA	United_Sta
Mr.	Alize	J.	Ferrari	alize_ferrar PhD		School of Public Health	University of Queensland	Brisbane	Queensland	Australia	Queensland Centre for Mental Health Research	University of Queensland	Brisbane	Queensland	Australia	Institute fo University i Seattle	Washington	United_Sta
Dr.	Michael		Kutz	kutz13@uv BS		Institute for Health Metrics and Evaluation	University of Washington											
Dr.	Janni		Leung	j.leung1@u PhD		School of Public Health	University of Queensland	Brisbane	QLD	Australia	Queensland Centre for Mental Health Research	University of Washington	Seattle	WA	United_States			
Dr.	Damian		Santomauro	damians@u PhD		School of Public Health	University of Queensland	Brisbane	Queensland	Australia	Queensland Centre for Mental Health Research	University of Washington	Seattle	WA	United_Sta			
Dr.	Haidong		Wang	haidong@u PhD		Institute for Health Metrics and Evaluation	University of Washington	Seattle	Washington	United_States								
Prof.	Harvey	A.	Whiteford	h.whiteford PhD		School of Public Health	University of Queensland	Brisbane	Queensland	Australia	Queensland Centre for Mental Health Research	University of Queensland	Brisbane	Queensland	Australia	Institute fo University i Seattle	United_Sta	
Mr.	Amanuel		Alemu	amanuel.at MPH		School of Public Health	University of Queensland	Brisbane	QLD	Australia								
Prof.	Foad		Abajobir	foad.abdall MD		Department of Neurology	Cairo University	Cairo		Egypt								
Mr.	Haftom		Niguse	hafniguse@MS			Mekelle University	Mekelle	Tigray	Ethiopia								
Dr.	Laith	J.	Abu-Raddad	lja2002@q PhD		Infectious Disease Epidemiology Group	Weill Cornell Medical College in Qatar	Doha		Qatar								
Dr.	Aliasghar		Ahmad Kiadaliri	aliasghar.al PhD		Department of Clinical Sciences Lund, Orthopedics, Clinical Epidemiology Unit	Lund University	Lund		Sweden								
Dr.	Alireza		Ahmadi	ahmadiar1 PhD			Kermanshah University of Medical Sciences	Kermanshah		Iran	Stockholm		Sweden					
Mrs.	Kedir	Yimam	Ahmed	kediryimam MPH			Debre Markos University	Debre Mari	Amhara	Ethiopia								
Mr.	Muktar	Beshir	Ahmed	muktar27@ MPH		College of Health Sciences, Department of Epidemiology, ICT and e-Learning Coordinator	Jimma University	Jimma	Oromiya	Ethiopia								
Dr.	Faris	Hasan	Al Lami	farislami@ PhD			Baghdad College of Medicine	Baghdad	Baghdad	Iraq								
Dr.	Khurshid		Alam	khurshid.b PhD		Murdoch Childrens Research Institute	The University of Melbourne	Parkville	Victoria	Australia	The University of Melbourne	Melbourne VIC	Australia	The Univer Sydney	NSW	Australia		
Ms.	Deena		Alasfoor	deena.alasf MSc			Ministry of Health	Al Khuwair	Muscat	Oman								
Dr.	Reza		Alizadeh-Navaei	reza_nava@ PhD		Gastrointestinal Cancer Research Center	Mazandaran University of Medical Sciences	Sari	Mazandaran	Iran								
Dr.	Juma	M	Alkaabi	J.KAABI@U FRCP			United Arab Emirates University	ALAIN UAE Abu Zaby	Al-Ain City	united_arab_emirates								
Prof.	Fatma		Al-Maskari	fatma.am@ PhD		College of Medicine & Health Sciences	United Arab Emirates University	Al-Ain City		united_arab_emirates								
Dr.	Rajaa		Al-Raddadi	saudiresear PhD			King Saud University	Jeddah	Makkah	Saudi_Arabia								
Dr.	Khalid	A.	Altrikawi	kaltirkawi@ MD			King Saud University	Riyadh		Saudi_Arabia								
Dr.	Nahla		Anber	nahlaanber PhD			Mansoura University	Mansoura		Egypt								
Dr.	Hossein		Ansari	haepidemii PhD		Health Promotion Research Center, Department of Epidemiology and Biostatistics	Zahedan University of Medical Sciences	Zahedan		Iran								
Mr.	Hamid		Asayesh	asayeshpsy PhD		Department of Medical Emergency, School of Paramedic	Qom University of Medical Sciences	Qom		Iran								
Dr.	Rana	Jawad	Asghar	jawad@alu MD			South Asian Public Health Forum	Islamabad		Pakistan								
Mr.	Tesfay		Mehari	tesfay_meh MS			Mekelle University	Mekelle	Tigray	Ethiopia								
Mr.	Tadesse		Awoke Ayele	tawoke7@IMS			University of Gondar	Gondar		Ethiopia								
Prof.	Till		Bärnighausen	tbaernig@I MD		Department of Global Health and Population, Harvard T H Chan School of Public Health	Harvard University	Boston	MA	United_States	Africa Health Research Institute	Mtubatuba	KwaZulu-N:South_Afri	Institute of Heidelberg Heidelberg		Germany		
Mr.	Umar		Bacha	umar.bach PhD		School of Health Sciences	University of Management and Technology	Lahore	Punjab	Pakistan								
Dr.	Aleksandra		Barac	aleksandral PhD		Faculty of Medicine	University of Belgrade	Belgrade	Belgrade	Serbia								
Dr.	Suzanne	L	Barker-Collo	s.barker-co PhD		School of Psychology	University of Auckland	Auckland		new_zealand								
Prof.	Bernhard	T	Baune	bernhard.b PhD		School of Medicine	University of Adelaide	Adelaide	South Australia	Australia								
Prof.	Shahrazad		Bazargan-Hejazi	shahrzadba PhD		College of Medicine	Charles R. Drew University of Medicine and Science	Los Angeles: CA		United_States	David Geffen School of Medicine	University of California at Los Angeles	Los Angeles: CA	United_States				
Dr.	Neeraj		Bedi	drnbedi@y MD			College of Public Health and Tropical Medicine	Jazan		Saudi_Arabia								
Dr.	Isabela	M	Bensenor	isabenseno PhD			University of São Paulo	São Paulo	São Paulo	Brazil								
Dr.	Adugnaw		Berhane	adugnawm PhD		College of Health Sciences	Debre Berhan University	Debre Berhan		Ethiopia								
Mr.	Addisu	Shunu	Beyene	addisushun MPH		College of Health and Medical Science	Haramaya University	Harar	Oromia	Ethiopia								

Dr. Mr.	Zulfiqar Dube	A. Jara	Bhutta Boneya	zulfiqar.bh.jaradube@MPH	Centre of Excellence in Women and Child Health Department of Public Health	Aga Khan University Debre Markos University	Karachi Debre Mar/Amhara	Pakistan Ethiopia	Centre for Global Child Health	The Hospital for Sick Children	Toronto	ON	Canada
Dr. Prof. Dr.	Rohan Nicholas Zahid	J. K. A	Borschmann Breitborde Butt	rohan.bors breitborde. PhD zabutt3@y		The University of Melbourne The Ohio State University Al Shifa Trust Eye Hospital	Melbourne VIC Columbus OH Rawalpindi Punjab	Australia United_States Pakistan		Murdoch Childrens Research Institute	Melbourne VIC		Australia
Dr. Mrs. Dr.	Ferrán Liliانا Hadi	G	CatalÀj-LÀpez Ciobanu Danawi	ferran_cats PhD liliانا.ciobi:MS hdanawi@:PhD	Department of Medicine School of Medicine	University of Valencia/INCLIVA Health Research Institute and CIBERSAM University of Adelaide Walden University	Valencia Adelaide South Australia Minneapolis Minnesota	Spain Australia United_States	Clinical Epidemiology Program	Ottawa Hospital Research Institute	Ottawa		Canada
Dr.	Amare		Deribew	amare.deri PhD	Nuffield Department of Medicine Department of Community Medicine, Faculty of Medicine	University of Oxford	Oxford	United_Kingdom		KEMRI-Wellcome Trust Research Programme	Kilifi		Kenya
Dr.	Samath	D.	Dharmaratne	samath62@MD		University of Peradeniya	Peradeniya	Sri_Lanka					
Prof. Mr.	Kerrie Aman	E Yesuf	Doyle Endries	auntykerrie PhD amanyesuf.MPH		RMIT University Arba Minch University	Bundoora VIC Arba Minch SNNPR	Australia Ethiopia		Australian National University	Canberra	ACT	Australia
Dr. Prof.	Emerito Jose Andr�	Aquino	Faraon Faro	eafaraon@ MD andre.faro. PhD	College of Public Health	University of the Philippines Manila Federal University of Sergipe	Manila Aracaju Sergipe	Philippines Brazil		Department of Health	Manila		Philippines
Dr. Mr.	Maryam Wubalem	S	Farvid Fekadu	mfarvid@h PhD wubfek@gi:MS	Department of Nutrition, Harvard T.H. Chan School of Public Health	Harvard University Bahir Dar University	Boston MA Bahir Dar	United_States Ethiopia		Harvard/MGH Center on Genomics, Vulnerable Populations, and Health Disparities, Mongan Institute for Health Policy, Massachusetts General Hospital	Boston	MA	United_States
Dr. Dr. Mr. Dr.	Seyed-Mohammad Florian Tsegaye Ababi	Tewelde Zergaw	Fereshtehnejad Fischer Gebrehiwot Giref	sm.feresht:PhD f.fischer@u:PhD tsegaye.tev MPH ababi_z@y PhD	Department of Neurobiology, Care Sciences and Society (NVS) School of Public Health	Karolinska Institutet Bielefeld University Jimma University Addis Ababa University	Stockholm Bielefeld North Rhine-Westphalia Jimma Oromia Addis Ababa	Sweden Germany Ethiopia Ethiopia					
Mr.	Melkamu Dedefo		Gishu	melkyed@t:MS		Haramaya University	Dire Dawa	Ethiopia		Kersa Health and Demographic Surveillance System	Harar		Ethiopia
Dr. Mr. Prof. Mr.	Alessandra Tesfa Randah Mitiku	Carvalho Dejenie Ribhi Teshome	Goulart Habtewold Hamadeh Hambisa	agoulart@t PhD tesfadej20k:MS randah@ag:DPHl mitikkuuf@MPH	College of Health and Medical Sciences	Center for Clinical and Epidemiological Research Center- Hospital Universitario- University of S�o Paulo University of Groningen Arabian Gulf University Haramaya University Hamdan Bin Mohammed Smart University	S�o Paulo S�o Paulo Groningen Manama Harar Harari	Brazil Netherlands Bahrain Ethiopia		Center of Check of Hospital Sirio Libanes Debre Berhan University	S�o Paulo S�o Paulo Debre Berhan		Brazil Ethiopia
Dr.	Samer		Hamidi	s.hamidi@l DrPH		Parc Sanitari Sant Joan de D��u - CIBERSAM	Dubai	united_arab_emirates					
Dr.	Josep Maria		Haro	27652jha@MD			Sant Boi de Llobregat (Barcelon)	Spain					
Dr.	Mohammad Sadegh		Hassanvand	hassanvanc PhD	Center for Air Pollution Research, Institute for Environmental Research		Tehran	Iran					
Dr. Prof. Dr. Dr.	Nobuyuki Mohamed Hsiang Abdullatif		Horita Hsairi Huang Husseinii	horitano@h MD mohamed.i MD lhuang@cl MD abdullatifh:PHD	Department of Pulmonology Department of Epidemiology Institute of Community and Public Health	Tehran University of Medical Sciences Yokohama City University Graduate School of Medicine Salah Azaiz Institute Cambridge Health Alliance Birzeit University	Yokohama Kanagawa Tunis Tunis Cambridge MA Birzeit	Japan Tunisia United_States Palestine					
Prof.	Mihajlo	B	Jakovljevic	sidartagoth PhD	Faculty of Medical Sciences	University of Kragujevac	Kragujevac	Central Serbia - Sun	The Center for Health Trends and Forecasts	Institute for Health Metrics and Evaluation (IHME) The University of Washington	Seattle	Washington	United_States
Dr.	Spencer	Lewis	James	spencj@gm MD		Denver Health/University of Colorado	Denver	CO	United_States				
Prof.	Jost	B	Jonas	Jost.Jonas@MD	Department of Ophthalmology, Medical Faculty Mannheim	Ruprecht-Karls-University Heidelberg	Mannheim	Germany	Germany				
Dr.	Amir		Kasaean	amir_kasae PhD	Hematology-Oncology and Stem Cell Transplantation Research Center Department of Community Medicine, Public Health and Family Medicine	Tehran University of Medical Sciences Jordan University of Science and Technology	Tehran Irbid	Tehran Irbid Punjab	Iran Jordan Pakistan	Endocrinology and Metabolism Population Sciences Institute	Tehran	Tehran	Iran
Prof. Dr. Dr.	Yousef Ejaz Abdullah	Saleh Ahmad Tawfi�	Khader Khan Khoja	yskheader@ ScD ejaz@hsa.e MD atkhoja@g:MD		Health Services Academy Mohammed Ibn Saudi University Iranian Ministry of Health and Medical Education	Irbid Islamabad Riyadh	Saudi Arabia	Saudi_Arabia	Non-Communicable Diseases Research Center	Tehran		United_States
Dr.	Ardeshir		Khosravi	ardeshir.kh PhD			Tehran	Iran			Tehran		Iran
Dr. Prof. Dr.	Jagdish Daniel Yun Jin		Khubchandani Kim Kim	jkhubchanc PhD daniel.kim@DrPH neurokim7:PHD	Department of Nutrition and Health Science Department of Health Sciences Faculty of Chinese Medicine	Ball State University Northeastern University Southern University College	Muncie Boston Skudai	Indiana Massachusetts Johor	United_States United_States Malaysia				
Dr.	Yoshihiro		Kokubo	ykokubo@l PhD	Department of Preventive Cardiology	National Cerebral and Cardiovascular Center Parc Sanitari Sant Joan de Deu (CIBERSAM)	Suita Osaka	Japan					
Dr.	Ai		Koyanagi	a.koyanagi@MD	Research and Development Unit		Barcelona	Spain					

Prof.	Barthelemy		Kuate Defo	Barthelemy PhD	Department of Social and Preventive Medicine, School of Public Health	University of Montreal	Montreal	Quebec	Canada	Department of Demography and Public Health Research Institute	University of Montreal	Montreal	Canada
	Heidi	J.	Larson	Heidi.Larso PhD	Department of Infectious Disease Epidemiology	London School of Hygiene & Tropical Medicine	London		United_Kingdom	Institute for Health Metrics and Evaluation	University of Washington	Seattle WA	United_States
Dr.	Asma	Abdul	Latif	asmaSlatif@PhD	Department of Zoology	Lahore College for Women University	Lahore	Punjab	Pakistan				
Dr.	Paul	H	Lee	paul.h.lee@PhD		Hong Kong Polytechnic University	Hong Kong		China				
Mr.	Cheru	Tesema	Leshargie	chertesem@MPH		Debre Markos University	Debre Mar	Amhara	Ethiopia				
Dr.	Ricky		Leung	rleung@altPhD		State University of New York, Albany	Rensselaer NY		United_States				
Dr.	Loon-Tzian		Lo	loonlo@un MD		UnionHealth Associates, LLC	St. Louis MO		United_States		Alton Mental Health Center	Alton IL	United_States
Prof.	Raimundas		Lunevicius	rlunevichusPhD		Aintree University Hospital National Health Service Foundation Trust	Liverpool		United_Kingdom	School of Medicine	University of Liverpool	Liverpool	United_Kingdom
Dr.	Hassan		Magdy Abd El Razek	farreha201 MB BCH		Mansoura Faculty of Medicine	Mansoura		Egypt				
Dr.	Mohammed		Magdy Abd El Razek	hoda_734@MB BCH	Aswan University Hospital	Aswan Faculty of Medicine	Aswan		Egypt				
Prof.	Reza		Majdzadeh	rezamajd@PhD	Knowledge Utilization Research Center and Community Based Participatory Research Center	Tehran University of Medical Sciences	Tehran		Iran				
Prof.	Azeem		Majeed	a.majeed@MD	Department of Primary Care & Public Health	Imperial College London	London	England	United_Kingdom	Digestive Diseases Research Institute	Tehran University of Medical Sciences	Tehran	Iran
Prof.	Reza		Malekzadeh	malek@tur MD	Digestive Diseases Research Institute	Tehran University of Medical Sciences	Tehran		Iran				
Dr.	Jose		Martinez-Raga	martinezra@PhD		Hospital Universitario Doctor Peset	Valencia	Valencia	Spain		CEU Cardinal Herrera University	Moncada Valencia	Spain
Dr.	Habibolah		Masoudi Farid	Habib.mas@MD		State Welfare Organisation	Tehran	TEHRAN	Iran				
Mr.	Mohsen		Mazidi	Mazidi_ns@PhD	Key State Laboratory of Molecular Developmental Biology, Institute of Genetics and Developmental Biology	Chinese Academy of Sciences	Beijing		China				
Prof.	John	J	McGrath	j.mcgrath@PhD	Queensland Centre for Mental Health Research	The Park Centre for Mental Health	Wacol	Queensland	Australia	Queensland Brain Institute	University of Queensland	St Lucia QLD	Australia
Prof.	Ziad	A.	Memish	zmemish@ MD		Saudi Ministry of Health	Riyadh		Saudi_Arabia	College of Medicine	Afaisal University	Riyadh	National Ce Aarhus Uni Aarhus
Dr.	Walter		Mendoza	mendoza@MD		United Nations Population Fund	Lima		Peru				
Mr.	Melkamu	Merid	Mengesha	melkamum@MPH	College of Health and Medical Sciences	Haramaya University	Harar	Kebele 18	Ethiopia				
Mr.	Mubarek	Abera	Mengstie	abmubarek@PhD Fellow	Center for Translation Research and Implementation Science, National Heart, Lung, and Blood Institute	Jimma University	Jimma	Oromia	Ethiopia				
Mr.	George Haftay	A. Berhane	Mensah Mezgebe	george.mei MD haftay.berf@MS		National Institutes of Health	Bethesda MD		United_States				
Dr.	Ted	R	Miller	miller@pir@PhD		Mekelle University	Mekelle	Tigray	Ethiopia				
Prof.	Philip	B.	Mitchell	phil.mitche MD		Pacific Institute for Research & Evaluation	Calverton MD		United_States	Centre for Population Health	Curtin University	Perth WA	Australia
Dr.	Alireza		Mohammadi	ar.mohamr PhD	Neuroscience Research Center	University of New South Wales	Kensington New South Wales		Australia				
Dr.	Shafiu		Mohammed	shafiu@PhD	Health Systems and Policy Research Unit Center for Research on Population and Health, Faculty of Health Sciences	Baqiyatallah University of Medical Science	Tehran Zaria	Kaduna	Nigeria	Institute of Public Health	Heidelberg University	Heidelberg Baden Wue	Germany
Prof.	Carla	Makhlouf	Obermeyer	cm39@aub@DSc	Center for Health Research	American University of Beirut	Beirut		Lebanon				
Dr.	Felix	Akpojene	Ogbo	felgbo@yal@MPH		Western Sydney University	Sydney	New South Wales	Australia				
Dr.	Elizabeth Christina		Palomares Castillo Papachristou	palomares@PhD christina.p@PhD		Ministry of Health	Mexico City	Mexico city	Mexico	Universidad Nacional Autónoma de México (UNAM)		Mexico City	Mexico city Mexico
Prof.	Scott	B.	Patten	patten@uc PhD	Department of Community Health Sciences	University of Calgary	Calgary	Alberta	Canada				
Prof.	George	C.	Patton	george.pat@MD	Murdoch Childrens Research Institute, Department of Paediatrics	University of Melbourne	Melbourne	Victoria	Australia				
Prof.	David	M.	Pereira	PHD	REQUIMTE/LAQV, Laboratório de Farmacognosia, Departamento de Química, Faculdade de Farmácia	Universidade do Porto	Porto		Portugal				
Dr.	Aslam		Pervaz	aslam_perv@MHA		Postgraduate Medical Institute	Lahore	Punjab	Pakistan				
Prof.	Michael Farshad	Robert	Phillips Pourmalek	mphillips@PhD farshad.poi@PhD		Shanghai Jiao Tong University School of Medicine	Shanghai		China		Emory University	Atlanta Georgia	United_States
Dr.	Mostafa Amir		Qorbani Radfar	mqorbani1 PhD aradfar@at@MD	Non-Communicable Diseases Research Center	University of British Columbia	Vancouver	British Columbia	Canada				
Dr.	Anwar		Rafay	rafay.aku.e@MS		Alborz University of Medical Sciences	Karaj		Iran				
Prof.	Vafa		Rahimi-Movaghar	v_rahimi@MD	Sina Trauma and Surgery Research Center	A T Still University	Kirkville MO		United_States				
Mr.	Rajesh Kumar		Rai	rajesh.jips2@MPH		Contech International Health Consultants	Lahore	Punjab	Pakistan		Contech School of Public Health	Lahore Punjab	Pakistan
Dr.	David	Laith	Rawaf	drawaf@ic.MD, MBBS	WHO Collaborating Centre	Tehran University of Medical Sciences	Tehran	Tehran	Iran				
Prof.	Salman		Rawaf	s.rawaf@in@MD		Society for Health and Demographic Surveillance	Suri	West Bengal	India		North Hampshire Hospitals	Basingstroke	United_Kingdom
Prof.	Amany	H	Refaat	amany.refa@MD		Imperial College London	London		United_Kingdom				
Dr.	Satar		Rezaei	satar.rezae@PhD	School of Public Health	Imperial College London	London		United_Kingdom				
Dr.	Mohammad Sadegh		Rezai	drmsrezai@MD		Walden University	Minneapolis MN		United_States		Suez Canal University	Ismailia Ismailia	Egypt
						Kermanshah University of Medical Sciences	Kermanshah		Iran				
						Mazandaran University of Medical Sciences	Sari	Mazandaran	Iran				

Dr.	Gholamreza		Roshandel	roshandel_PhD	Golestan Research Center of Gastroenterology and Hepatology	Golestan University of Medical Sciences	Gorgan		Iran	Digestive Diseases Research Institute	Tehran University of Medical Sciences	Tehran	Iran						
Dr.	Mahdi		Safdarian	mahdisafde MD		Sina Trauma & Surgery Research Center	Tehran		Iran										
Dr.	Mahdi		Safiabadi	fcf.drsafiat MD	Student Research Committee Managerial Epidemiology Research Center, Department of Public Health, School of Nursing and Midwifery	Baqiyatallah University of Medical Sciences	Tehran		Iran										
Dr.	Saeid		Safiri	saeidsafiri@PhD		Maragheh University of Medical Sciences	Maragheh		Iran										
Dr.	Rajesh		Sagar	rsagar29@iMD		All India Institute of Medical Sciences	New Delhi	Delhi	India										
Dr.	Mohammad Ali		Sahraian	sahraian13 MD	MS Research Center, Neuroscience Institute	Tehran University of Medical Sciences	Tehran		Iran										
Prof. Dr.	Payman Abdallah	M.	Salamati Samy	psalamati@MD asamy.ku@PhD	Sina Trauma and Surgery Research Center	Tehran University of Medical Sciences	Tehran	Tehran	Iran										
Prof. Dr.	Benn Mete Soraya	I	Sartorius Saylan Seedat	sartorius@ PhD msaylan@i sseedat@si PhD	Public Health Medicine, School of Nursing and Public Health	University of KwaZulu-Natal	Durban		South_Africa	UKZN Gastrointestinal Cancer Research Centre	South African Medical Research Council (SAMRC)	Durban	South_Africa	Lawrence	Kansas	United_States			
Prof. Dr.	Sadaf Masood	G	Ali	sepanlou@ PhD masoodalis MD	Digestive Diseases Research Institute	Tehran University of Medical Sciences	Tehran	Tehran	Iran										
Mr.	Morteza		Shamsizadeh	mshamsizadeh MPH	Department of Medical Surgical Nursing, School of Nursing and Midwifery	Hamadan University of Medical Sciences	Hamadan		Iran										
Dr.	Diego	Augusto	Santos	silva diegoaugus PhD		Federal University of Santa Catarina	Florianopolis	SANTA CATARINA	Brazil										
Dr. Dr.	Jasvinder Badr	A. H	A	Singh Sobaih	jasvinder.m MD bsobaih@k MD	Medical Center king Saud university	Birmingham	Alabama	United_States					Riyadh	Saudi_Arabia				
Dr.	Chandrashekhar		T.	Sreeramareddy	MD	Department of Community Medicine	International	Kuala Lumpur,	Malaysia										
Prof.	Dan	J		Stein	dan.stein@ PhD	Department of Psychiatry	University of Cape Town	Cape Town	Western Province	South_Africa				South African Medical Research Council Unit on Anxiety & Stress Disorders	Cape Town	Western Cape	South_Africa		
Dr.	Rizwan			Suliankatchi Abdulkad sarizwan19 MD		Ministry of Health, Kingdom of Saudi Arabia	Riyadh		Saudi_Arabia										
Prof.	Bryan	L		Sykes	blysykes@ur PhD	Departments of Criminology, Law & Society, Sociology, and Public Health	University of California, Irvine	Irvine	CA	United_States									
Prof. Dr.	Rafael Muawiyiyah			Tabarés-Seisdedos Sufiyan	rafael.tabar@ PhD MD	Department of Medicine	University of Valencia, INCLIVA Health Research Institute and CIBERSAM	Valencia	Valencia	Spain									
Dr.	Karen	M.		Tabb	ktabb@illini PhD	School of Social Work Preventive Medicine and Public Health Research Center	University of Illinois at Urbana-Champaign	Champaign	Illinois	United_States									
Dr. Dr.	Arash Mohamad-Hani Abdullah			Tehrani-Banihashemi Temsah Terkawi	tehranib@i PhD Temsah1@ MD asterkawi@ MD	Department of Anesthesiology Institute of Public Health, Faculty of Health Sciences	Iran University of Medical Sciences	Tehran	Tehran	Iran									
Dr. Dr.	Roman Kingsley Olalekan	Nnanna A		Topor-Madry Ukwaja Uthman	roman.topr@ PhD ukwajaking MD olalekan.ut PhD	Department of Internal Medicine Warwick Medical School	King Saud University	Riyadh	United_States	Department of Anesthesiology	King Fahad Medical City	Riyadh	Saudi_Arab	Outcomes	Cleveland	Cleveland	OH	United_States	
Dr. Dr.	Roman Kingsley Olalekan	Nnanna A		Topor-Madry Ukwaja Uthman	roman.topr@ PhD ukwajaking MD olalekan.ut PhD	Department of Internal Medicine Warwick Medical School	Jagiellonian University Medical College	Kraków	Poland	Faculty of Health Sciences	Wrocław Medical University	Wrocław	Poland						
Prof. Mr.	Stein Emil Tolassa			Vollset Wakayo	vollset@uil DrPH wakayotola MS	Center for Disease Burden	Norwegian Institute of Public Health	Bergen	NA	Norway	Department of Global Public Health and Primary Care	University of Bergen	Bergen	Norway	Institute fo	University of Seattle	WA	United_States	
Dr.	Yuan-Pang			Wang	gnap_inbo@ PhD	Competence Center Mortality-Follow-Up of the German National Cohort	University of São Paulo Medical School	São Paulo	São Paulo	Brazil									
Dr.	Andrea			Werdecker	andrea.wer@ PhD		Federal Institute for Population Research	Wiesbaden	Hessen	Germany									
Dr. Mr. Mr. Mr.	Ronny Abdulhalik Mohsen Hassen		Hamid	Westerman Workicho Yaghoubi Yimam	ronny.west@ PhD abdulhalik MPH mohsen.ya@ MA abdulhamid MPH	Department of Biostatistics, School of Public Health	Iran University of Medical Sciences	Tehran	Tehran	Iran	German National Cohort Consortium	Heidelberg	Baden-Wur	Germany	Ghent	Belgium	Tehran	Tehran	Iran
Mr. Prof.	Naohiro Mustafa	Z		Yonemoto Younis	nyonemotc MPH younis68@ DrPH	Department of Biostatistics, School of Public Health	Kyoto University	Kyoto	Japan	Department of Global Public Health and Primary Care	University of Bergen	Bergen	Norway	Institute fo	University of Seattle	WA	United_States		
Prof. Prof.	Chuanhua Maysaa	El Sayed		Yu Zaki	yuchua@w PhD may_s65@ PhD	Department of Epidemiology and Biostatistics, School of Public Health Faculty of Medicine	Wuhan University	Wuhan	Hubei	China	Global Health Institute	Wuhan University	Wuhan	Hubei	China				
Dr. Prof. Prof.	Aisha Theo Mohsen	O		Jumaan Vos Naghavi	aishaoj@gr PhD tvos@uw.e PhD nagham@u PhD	Institute for Health Metrics and Evaluation	Independent Consultant	Seattle	Washington	United_States									
Prof.	Simon	I.		Hay	sihay@uw. DSc	Oxford Big Data Institute, Li Ka Shing Centre for Health Information and Discovery	University of Oxford	Oxford	United_Kingdom	Institute for Health Metrics and Evaluation	University of Washington	Seattle	Washington	United_States					
Dr. Prof.	Nicholas Christopher	J	L	Kassebaum Murray	nickjk@uw MD cjlm@uw.e DPhil	Institute for Health Metrics and Evaluation	University of Washington	Seattle	Washington	United_States	Department of Anesthesiology & Pain Medicine	Seattle Children's Hospital	Seattle	Washington	United_States				

