



Burden of lower respiratory infections in the Eastern Mediterranean Region between 1990 and 2015: findings from the Global Burden of Disease 2015 study

GBD 2015 Eastern Mediterranean Region LRI Collaborators · Ali H. Mokdad¹

Received: 1 May 2017 / Revised: 8 June 2017 / Accepted: 28 June 2017
© The Author(s) 2017. This article is an open access publication

Abstract

Objectives We used data from the Global Burden of Disease 2015 study (GBD) to calculate the burden of lower respiratory infections (LRIs) in the 22 countries of the Eastern Mediterranean Region (EMR) from 1990 to 2015.

Methods We conducted a systematic analysis of mortality and morbidity data for LRI and its specific etiologic factors, including pneumococcus, *Haemophilus influenzae* type b, Respiratory syncytial virus, and influenza virus. We used modeling methods to estimate incidence, deaths, and disability-adjusted life-years (DALYs). We calculated burden attributable to known risk factors for LRI.

Results In 2015, LRIs were the fourth-leading cause of DALYs, causing 11,098,243 (95% UI 9,857,095–12,396,566) DALYs and 191,114 (95% UI 170,934–210,705) deaths. The LRI DALY rates were higher than global estimates in 2015. The highest and lowest age-standardized rates of DALYs were observed in Somalia and Lebanon, respectively. Undernutrition in

childhood and ambient particulate matter air pollution in the elderly were the main risk factors.

Conclusions Our findings call for public health strategies to reduce the level of risk factors in each age group, especially vulnerable child and elderly populations.

Keywords Lower respiratory infection · Incidence · Mortality · DALY · Eastern Mediterranean Region

Introduction

Lower respiratory infections (LRIs) are one of the leading causes of death and disability-adjusted life-years (DALYs) worldwide, especially among children under 5 (Kassebaum et al. 2016; Wang et al. 2016). Since 1990, LRI has been the closest competitor of cardiovascular diseases as the top leading cause of DALYs in the Eastern Mediterranean Region (EMR) and the Arab world (Mokdad et al. 2014, 2016). Several factors, such as poverty, indoor and outdoor air pollution, malnutrition, smoking, chronic lung diseases, and delayed and inappropriate case management contribute to the high burden of LRI (Hadi 2003). Population-based measures of morbidity and mortality of LRI are scarce. Most of the available data are limited to children and are based on modeling approaches. Even fewer data are available for causative agents of LRI (Kovacs et al. 2015).

Rudan et al. estimated an incidence of 0.22 episodes of community-acquired pneumonia per child-year in children under 5 in low- and middle-income countries in 2010 (Rudan et al. 2013). The Child Health Epidemiology Reference Group (CHERG) estimated 0.26 episodes per child-year for the world and 0.28 for the World Health Organization's (WHO) Eastern Mediterranean Region (EMR).

Corresponding author: Ali H. Mokdad.

This article is part of the supplement “The state of health in the Eastern Mediterranean Region, 1990–2015”.

The members of GBD 2015 Eastern Mediterranean Region LRI are listed at the end of the article.

Electronic supplementary material The online version of this article (doi:10.1007/s00038-017-1007-0) contains supplementary material, which is available to authorized users.

✉ Ali H. Mokdad
mokdaa@uw.edu

¹ Institute for Health Metrics and Evaluation, 2301 5th Avenue, Suite 600, Seattle, WA 98121, USA

These estimates translate to about 20 million cases of childhood pneumonia each year in the EMR, with approximately 10% of cases requiring hospitalization (Rudan et al. 2008). More than 99% of all LRI deaths occur in low- and middle-income countries. Although about 62% of children with severe LRI reach hospitals, more than 80% of all childhood LRI deaths take place outside the hospital setting (Nair et al. 2013; Tong 2013). Despite high levels of morbidity and mortality, the decreasing trend in LRI mortality rates has contributed to increasing life expectancy worldwide (Wang et al. 2016). Improvements in nutritional status (less malnutrition through attention to childhood and maternal nutritional status), increased uptake of vaccines such as immunization against *Streptococcus pneumoniae* (pneumococcus) and *Haemophilus influenzae* type b (Hib) in children and high-risk populations in EMRs, and improved access to antibiotics and supportive care have decreased the incidence and fatality of LRI in many countries (Williams and Shah 2012; Tong 2013).

In this study, we report findings from the Global Burden of Diseases, Injuries, and Risk Factors Study 2015 (GBD 2015) on LRI in the 22 countries of the EMR between 1990 and 2015. We describe the burden of LRI and a subset of specific etiologic pathogens, including pneumococcus, Hib, respiratory syncytial virus (RSV), and influenza virus, based on deaths, years of life lost (YLLs), years lived with disability (YLDs), and disability-adjusted life-years (DALYs).

Methods

GBD 2015 covers 195 countries, 21 regions, and seven super-regions from 1990 to 2015 for 315 diseases and injuries and 79 risk factors. A counterfactual approach was used for estimating LRI etiologies, one of the methodological differences compared to GBD 2010. Detailed descriptions of the methodological approach for GBD 2015 have been published elsewhere (Forouzanfar et al. 2016; Kassebaum et al. 2016; Vos et al. 2016; Wang et al. 2016).

The EMR consists of 22 countries with different levels of gross national income (GNI) per capita. The low-income countries (LICs) are Afghanistan, Djibouti, Somalia, and Yemen. The middle-income countries (MICs) are Egypt, Iran, Iraq, Jordan, Lebanon, Libya, Morocco, Pakistan, Palestine, Sudan, Syria, and Tunisia. The high-income countries (HICs) are Bahrain, Kuwait, Oman, Qatar, Saudi Arabia, and the United Arab Emirates (UAE).

All types of LRI (bronchitis, bronchiolitis, and pneumonia) were included in this study. We used the following International Classification of Diseases (ICD-10) codes (or their corresponding codes from earlier ICD versions) as

equivalent to LRI: J09–J15.8, J16–J16.9, J20–J21.9, P23–P23.9, and Z25.1. The ICD-10 codes for etiologic categories of LRI included J09–J11.89 and Z25.1 for Influenza virus, J12.1 for RSV, J13, J13.0, J15.3, J15.4, and J15.6 for pneumococcus, J14 and J14.0 for Hib, and J12, J12.0, J12.2–J12.9, J15–J15.9, J15.5, J15.7, J15.8, J16–J16.9, J20–J21.9, and P23–P23.9 for other LRI. We did not include tuberculosis in this study; it has been classified as a separate item in GBD.

An analysis of available data on all-cause mortality for all countries was undertaken (Wang et al. 2016). For pathogen-specific mortality rates of LRI, we used a counterfactual approach based on the epidemiological concept of attributable mortality. The change in LRI was estimated assuming the condition that a specific pathogen was not present. We adopted different approaches to estimate bacterial and viral causes based on the available data. For pneumococcal and Hib LRI, we estimated the causal fraction from vaccine efficacy trials data. For RSV and influenza, we relied on observational studies that measured causal fractions among hospital admissions for LRI (Vos et al. 2016; Wang et al. 2016). We estimated the causal fractions among cases by country, age, and sex. To account for the higher case-fatality rate of bacterial versus viral LRI, we applied a relative case-fatality differential based on hospital data that included cases coded to the specific pneumonia causes. Our mortality estimates were used to calculate cause-specific YLLs for each age, sex, location, and calendar year.

To estimate LRI-related morbidity, a systematic review of studies on epidemiological indicators of LRI was done as part of the GBD standard methodology. We used 197 sources of data from EMR countries (46 for non-fatal outcomes and others for cause of death) which contained data on LRI. A list of all data sources is available on the Institute for Health Metrics and Evaluation's website. (Institute for Health Metrics and Evaluation) A series of Bayesian meta-regression analyses through DisMod-MR 2.1 were used for disease modeling. Model-based epidemiological estimates in combination with disability weights were used to calculate cause-specific YLDs for each age, sex, location, and calendar year. DALYs were calculated through summation of YLLs and YLDs (Kassebaum et al. 2016; Vos et al. 2016).

We estimated burden of LRI attributable to childhood malnutrition (underweight, wasting, and stunting), non-exclusive breastfeeding, zinc deficiency, smoking, secondhand smoke exposure, household air pollution from solid fuels, and ambient air pollution as known LRI risk factors. Details on definitions and measures of age- and sex-specific relative risks of LRI for each of the risk factors are available elsewhere (Forouzanfar et al. 2016).

We report 95% uncertainty intervals (UI) for each estimate—such as rates or numbers of deaths or DALYs.

We estimated UIs by taking 1000 samples from the posterior distribution of each quantity and using the 25th- and 975th-ordered draws of the uncertainty distribution.

Results

LRI deaths decreased by 30.2% from 273,714 (95% UI 239,890–306,648) in 1990 to 191,114 (95% UI 170,934–210,705) in 2015 in EMR countries. In spite of the decrease, LRI is still the third-leading cause of death for all ages in 2015. The death rate per 100,000 was 74.0 (95% UI 64.8–82.9) in 1990 and decreased to 29.5 (95% UI 26.4–32.5) in 2015, a 60.1% reduction. The age-standardized death rate for LRI was 65.4 (95% UI 58.6–73.4) in 1990 and declined by 38.5% to 40.2 (95% UI 36.0–45.2) in 2015. In 2015, 4.8% (95% UI 4.3–5.2) of all deaths and 13.0% (95% UI 11.5–14.7) of under-5 deaths were due to LRI.

At the regional level, the highest death rates and numbers of deaths were among children under 5 years old, followed by those aged 65 years or older. However, there was considerable heterogeneity in the age pattern of LRI deaths between the countries of the EMR. In all high-income countries (Bahrain, Saudi Arabia, Kuwait, Oman, Qatar, and UAE), and some of the middle-income countries

(Iran, Lebanon, Libya, Morocco, Palestine, Syria, and Tunisia), the LRI death rates among older age groups were greater than the mortality rates among children under 5 years. Death rates were 31.5 per 100,000 (95% UI 27.3–35.4) in men compared to 27.4 per 100,000 in women (95% UI 23.9–31.2) in the region. The LRI death rate was not different in EMR countries in boys versus girls under 5 (2015).


Figure 1 shows the age-standardized death rates for LRI by EMR country in 2013. Somalia, Djibouti, and Afghanistan had the highest death rates in the region.

Among the etiologic causes, pneumococcus had the highest mortality rate (16.6 per 100,000, 95% UI 10.0–22.9), followed by respiratory syncytial virus (1.5 per 100,000, 95% UI 0.9–2.4), *Haemophilus influenzae* type b (1.1 per 100,000, 95% UI 0.0–2.3), and influenza virus (0.8 per 100,000, 95% UI 0.5–1.2). Figure 2 demonstrates the contribution of different etiologic causes to LRI deaths by age.

The rate of YLLs per 100,000 population decreased by 69.5% from 5579 (95% UI 4,831–6,320) in 1990–1702 (95% UI 1,510–1,902) in 2015. However, LRI was still the third-leading cause of YLLs in 2015.

e-Table 1 shows age-standardized incidence rates of LRI in males and females of the EMR countries. The highest incidence rate was observed in Afghanistan and the

Fig. 1 Age-standardized death rates of lower respiratory tract infections per 100,000 population in the countries of the Eastern Mediterranean Region, Global Burden of Disease study, 2015






Fig. 2 Number of lower respiratory tract infection deaths by etiologic causes in the Eastern Mediterranean Region, Global Burden of Disease study, 2015

lowest in Tunisia for both sexes. In all countries but Afghanistan, Bahrain, and Djibouti, the incidence rates were higher among men than women, with the highest male-to-female ratio in Iran and Libya.

The rate of YLDs per 100,000 decreased from 15.8 (95% UI 10.7–22.9) to 9.7 (95% UI 6.5–13.6) per 100,000 during 1990–2015.

Total DALYs from LRI decreased by 40.6%, from 20,746,747 (95% UI 17,954,899–23,444,142) in 1990 to 11,098,243 (95% UI 9,857,095–12,396,566) in 2015. LRI, which was the leading cause of DALYs in 1990, was ranked fourth in 2015, behind ischemic heart disease, neonatal preterm birth complications, and neonatal encephalopathy. LRI DALY rates were 5594 (95% UI 4851–6334) and 1712 (95% UI 1520–1912) DALYs per 100,000 population in 1990 and 2015, respectively, a 69.4% reduction. About 4.9% (95% UI 4.3–5.5) of DALYs in all ages and 12.2% (95% UI 10.8–13.8) of DALYs in children under 5 were due to LRI in 2015.

Age-standardized DALY rates were 3411 per 100,000 (95% UI 2,993–3,824) in 1990 compared to 1518 per 100,000 (95% UI 1357–1673) in 2015 in the EMR. Global age-standardized DALY rates for LRI were lower than those in the EMR: 3310 (95% UI 3033–3551) in 1990, and 1428 (95% UI 1330–1511) in 2013. There was a 56.9% reduction in age-standardized DALY rates at the global level, which was similar to the EMR (55.5% reduction) during 1990–2015. Figure 3 presents trends in age-standardized and all-ages LRI DALY rates between 1990 and 2015 globally and for the EMR.

Figure 4 shows age-standardized LRI DALY rates in the EMR countries. The highest rates were seen in Somalia,

Afghanistan, and Djibouti. LRI DALY rates did not differ significantly between males and females.

Figure 5 shows DALY rates for different etiologies of LRI in EMR countries in 2015. The largest variation in country-specific DALYs (highest to lowest ratio) was observed for Hib (2068.7) compared to 125.5 for pneumococcus, 109.4 for RSV, and 30.6 for influenza. The DALY rate for Hib was less than one DALY per 100,000 in UAE and Qatar, compared to 916.7 in Somalia and 458.5 in Afghanistan (Fig. 5).

Figure 6 shows the LRI DALY rates attributable to different risk factors in 2015. Childhood undernutrition, household air pollution from solid fuels, ambient particulate matter pollution, suboptimal breastfeeding, second-hand smoke, no handwashing with soap, and zinc deficiency were the most important risk factors for LRI in children under 5. Among those aged 70 years or older, ambient particulate matter pollution, household air pollution from solid fuels, smoking, no handwashing with soap, secondhand smoke, and alcohol use were the main risk factors.

Discussion

Our study showed a substantial burden of LRI in the EMR. Despite a decline in the burden of LRI since 1990, LRI remained the fourth-leading cause of DALYs and third-leading cause of death in the region in 2015. The all-ages burden of LRI in the EMR is considerably higher than global rates, and even the age-standardized burden is slightly higher. Our findings call for intensified efforts to

Fig. 3 Trends of age-standardized (AS) and all-age (AA) DALY rates of lower respiratory infections in the Eastern Mediterranean Region and the world, Global Burden of Disease study, 1990–2015





Fig. 4 Age-standardized disability-adjusted life-years (DALYs) rates of lower respiratory tract infections per 100,000 population in the countries of Eastern Mediterranean Region, Global Burden of Disease study, 2015



control and reduce the burden of LRI in the EMR, especially among low-income countries and low-resource settings. Moreover, since the current situation of health status and LRI varies widely between the EMR countries, suggested interventions to avert LRI deaths and disabilities should be specified at the national or even subnational level (Akseer et al. 2015).

The unstable political situation and wars in many of the EMR countries present a major challenge for controlling communicable diseases in the region (Haq et al. 2013). Countries with highest burden of LRI (Somalia, Afghanistan, and Djibouti) have been involved in armed conflicts and social unrest in recent decades. Conflict-affected children experience significant increases in deaths due to

diarrheal diseases, malnutrition, respiratory infections, and measles (Guha-Sapir and Panhuis 2004). These are due to different causes such as displacement of populations, breakdown of health and social services, and increased risk of disease transmission (Murray et al. 2008). Beyond improving social determinants that affect LRI, a combination of public health and clinical strategies is required to control them (World Health Organization and UNICEF 1986). The “Protect, Prevent, and Treat” framework provides a blueprint for such interventions (Qazi et al. 2015). The need for an integrated global action plan led to the launch of the Global Action Plan for the Prevention and Control of Pneumonia and Diarrhea (GAPPD), which can be used by EMR countries as well (Qazi et al. 2015).

Fig. 5 Disability-adjusted life-years (DALYs) rates for different etiologies of lower respiratory tract infections in the countries of the Eastern Mediterranean Region, Global Burden of Disease study, 2015





Fig. 6 Disability-adjusted life-years (DALYs) attributable to risk factors of lower respiratory tract infections, Eastern Mediterranean Region, Global Burden of Disease study, 2015

Immunization is one of the most effective public health strategies to prevent and reduce the burden of LRI, such as immunization against pneumococcal infection, Hib, and seasonal influenza. Immunization is highly effective at

reducing the burden of ill health from pneumococcal infection and Hib, but less so for influenza. The effectiveness of influenza vaccination depends on the match between the antigenic strains included in the vaccine and

the circulating strains in the community, and this can vary substantially from year to year. EMR countries are at different stages of adopting these vaccines. High-income EMR countries have introduced pneumococcal conjugate vaccine (PCV) to their national immunization programs. Moreover, all low-income countries in the region except Somalia have introduced PCV thanks to support from Gavi, the Vaccine Alliance. (Gavi 2017 the Vaccine Alliance) Somalia has issues with under-coverage of immunization, even for traditional vaccines such as DTP vaccine (diphtheria, tetanus, and pertussis). The majority of middle-income countries in the region (Egypt, Iran, Iraq, Jordan, Lebanon, Syria, and Tunisia), have yet to introduce PCV or are in the early stages of planning (Moradi-Lakeh and Esteghamati 2013; Sibak et al. 2015).

All EMR countries have introduced Hib vaccine to their immunization programs, and coverage of Hib3 in the EMR has increased from 13% in 2005 to 72% in 2014. However, coverage of Hib3 varies across EMR countries (World Health Organization 2017). Hib3 coverage was 75% or less in five countries in 2014: Somalia, Syria, Iraq, Pakistan, and Afghanistan (World Health Organization 2017). The coverage rate in Syria dropped from 80% in 2010 to 43% due to the ongoing war in the country (World Health Organization 2017). Iran was the last country in the region to introduce Hib vaccine in late 2014 and has not yet published coverage data. Economic sanctions against Iran had an impact on health policies and slowed down implementation of immunization programs and policies (Kheirandish et al. 2015; Massoumi and Koduri 2015).

There are different immunization policies against seasonal influenza virus in EMR countries. Although the impact of the influenza vaccine on prevention of LRI is not as clear as that of conjugate vaccines against pneumococcal infections and Hib, its effectiveness in protecting specific high-risk groups against severe influenza-associated disease and death is generally accepted (Voordouw et al. 2003; Nichol et al. 2007). Indeed, the policies for influenza vaccination are impacted by national capacities, resources, and epidemiological status (World Health Organization 2012). In the countries that have already established vaccination programs, a number of strategies such as public education and reminders for parents and immunization providers can increase immunization coverage (Williams et al. 2011).

Improving diet and nutritional status in the population groups that are most vulnerable to LRI, such as children and elderly people, can help prevent infection. Additionally, enhancing breastfeeding and encouraging zinc supplementation can reduce the burden of childhood LRI in the region (Roth et al. 2008, 2010; Shah et al. 2013). Moreover, based on our data, reducing indoor air pollution (mainly due to household use of solid fuels) and

outdoor air pollution (especially ambient particulate matter) contribute to reducing the burden of disease in all age groups (Renzetti et al. 2009; Rehfuess et al. 2009). Unfortunately, many countries of the region suffer from air pollution: while Afghanistan, Pakistan, Somalia, Sudan, and Yemen have considerable issues with household air pollution, more countries are affected by outdoor air pollution (Afghanistan, Egypt, Iran, Iraq, Lebanon, Libya, Pakistan, Saudi Arabia, Syria, and Tunisia) (Cohen et al. 2017; World Health Organization Eastern Mediterranean Region 2017). Smoking and exposure to secondhand smoke are other factors that influence LRI (Oberge et al. 2011). Most of the countries of the region (all except Somalia and Morocco) have ratified the Framework Convention on Tobacco Control (FCTC), which provides a baseline for tobacco control (Framework Convention Alliance Parties to the WHO 2017). However, the anti-smoking activities need to be boosted (Usmanova and Mokdad 2013) and expanded to shisha use, which is very common in the region (Moradi-Lakeh et al. 2015).

Access to adequate health care and appropriate case management (rapid diagnosis, access to antibiotics and supportive care) are important therapeutic interventions for LRI control (Bhutta et al. 2013). Although in general health care access has improved in EMR countries, there is still limited access to high-quality health care due to availability, accessibility, acceptability, and costs, especially for those who live in remote areas as well as disadvantaged groups (Kronfol 2012; Takian et al. 2013). The use of simple, standardized guidelines for diagnosis and treatment of LRI reduces complications and deaths. These guidelines should be made available to all care facilities including community health facilities and mobile units (Kronfol 2012; Takian et al. 2013). Unfortunately, many EMR countries have not implemented such evidence-based guidelines. Although the required expertise and local data for developing guidelines are not available in the majority of EMR countries, adaptation of international guidelines can be used as an alternative method to increase quality of care (Rashidian 2008). On the other hand, several countries in the region (such as Libya, Syria, and Yemen, as the most recent examples) have experienced considerable setbacks in access to health care because of war, social unrest, and mass displacement of populations (Dewachi et al. 2014; Burki 2015). Shortages in the provision of acute care, inpatient and outpatient clinical services as well as primary care services, and access to essential medicines are among the most urgent issues in such unstable conditions. Moreover, continuity of providing preventive and health promotion services is difficult during war and instability; several reports have demonstrated the effect of war on childhood malnutrition

and the inability to maintain a cold chain for vaccines during the wars in Afghanistan, Iraq, and Yemen (Assefa et al. 2001; Levy and Sidel 2016; Burki 2016). These are among the well-known LRI risk factors.

Surveillance systems are crucial for planning and managing LRI and identifying emerging infections (Breiman et al. 2013). In 2015, several outbreaks of LRI occurred in the EMR, for instance influenza A(H1N1) pdm09 in Libya, Kuwait, and Jordan; avian influenza A (H5N1) in Egypt; and Middle East respiratory syndrome coronavirus (MERS-CoV) in Saudi Arabia and Jordan (WHO Regional Office for the Eastern Mediterranean 2016). However, many EMR countries do not have adequate health information systems to monitor the situation of LRI. (World Health Organization and UNICEF 2013) Therefore, it is important to assist countries to set up such systems to identify and manage outbreaks, and monitor the trends and risk factors of LRI. It also helps to improve accuracy of estimates for EMR in the next rounds of the GBD study.

Our study has some limitations. First, there are not enough original data for all countries of the region. For instance, many countries did not have estimates on incidence or etiologic agents of LRI. However, we used modeling approaches to estimate the burden, even for countries with limited data. While we did not assess avian influenza, severe acute respiratory syndrome (SARS), and MERS-CoV (Beauté et al. 2014; Milne-Price et al. 2014) as separate etiologic agents, all of them were included in total numbers of incidence, deaths, and DALYs.

Conclusion

LRI is one of the leading causes of morbidity and mortality in the EMR. Efforts are urgently needed to prevent and control LRI in the region, especially in the low-income countries of the region. All countries should consider adopting vaccines against pneumococcus, if they have not already, to reduce the burden. To adequately address the LRI burden, a public health and health care systems approach is needed. A comprehensive plan that includes addressing known risk factors such as poor diet, smoking, and exposure to secondhand smoke; health systems to improve prevention and treatment of cases; and community programs to increase awareness and immunization uptake are needed. A reliable surveillance system is required to monitor trends of disease burden and evaluate new interventions.

Unstable economies, political instability, and unrest in the region are major barriers to improving health. Therefore, it is important for the international community to work to improve political stability in the EMR, and to strengthen support for low-income countries and disadvantaged groups, in order to reduce LRI morbidity and mortality in the region.

GBD 2015 Eastern Mediterranean Region LRI Collaborators: Maziar Moradi-Lakeh MD, Department of Community Medicine, Preventative Medicine and Public Health Research Center, Gastrointestinal and Liver Disease Research Center (GILDRC), Iran University of Medical Sciences, Tehran, Iran. Charbel El Bcheraoui, PhD, Institute for Health Metrics and Evaluation, University of Washington, Seattle, Washington, United States. Raghid Charara, MD, American University of Beirut, Beirut, Lebanon. Ibrahim Khalil, MD, Institute for Health Metrics and Evaluation, University of Washington, Seattle, Washington, United States. Ashkan Afshin, MD, Institute for Health Metrics and Evaluation, University of Washington, Seattle, Washington, United States. Nicholas J. Kassebaum, MD, Institute for Health Metrics and Evaluation, University of Washington, Seattle, Washington, United States; Department of Anesthesiology and Pain Medicine, Seattle Children's Hospital, Seattle, Washington, United States. Michael Collison, BS, Institute for Health Metrics and Evaluation, University of Washington, Seattle, Washington, United States. Farah Daoud, BA/BS, Institute for Health Metrics and Evaluation, University of Washington, Seattle, Washington, United States. Adrienne Chew, Institute for Health Metrics and Evaluation, University of Washington, Seattle, Washington, United States. Kristopher J. Krohn, BA, Institute for Health Metrics and Evaluation, University of Washington, Seattle, Washington, United States. Danny Colombara, PhD, Institute for Health Metrics and Evaluation, University of Washington, Seattle, Washington, United States. Rebecca Ehrenkrantz, MPH, Institute for Health Metrics and Evaluation, University of Washington, Seattle, Washington, United States. Kyle J. Foreman, PhD, Institute for Health Metrics and Evaluation, University of Washington, Seattle, Washington, United States; Imperial College London, London United Kingdom. Joseph Frostad, MPH, Institute for Health Metrics and Evaluation, University of Washington, Seattle, Washington, United States. William W. Godwin, BS, Institute for Health Metrics and Evaluation, University of Washington, Seattle, Washington, United States. Michael Kutz, BS, Institute for Health Metrics and Evaluation, University of Washington, Seattle, Washington, United States. Puja, C. Rao, MPH, Institute for Health Metrics and Evaluation, University of Washington, Seattle, Washington, United States. Robert Reiner, BA, Institute for Health Metrics and Evaluation, University of Washington, Seattle, Washington, United States. Christopher Troeger, MPH, Institute for Health Metrics and Evaluation, University of Washington, Seattle, Washington, United States. Haidong Wang, PhD, Institute for Health Metrics and Evaluation, University of Washington, Seattle, Washington, United States. Amanuel Alemu Abajobir, MPH, School of Public Health, University of Queensland, Brisbane, QLD, Australia. Kaja M. Abbas, PhD, Virginia Tech, Blacksburg, VA, United States. Semaw Ferede Abera, MSc, School of Public Health, College of Health Sciences, Mekelle University, Mekelle, Ethiopia; Food Security and Institute for Biological Chemistry and Nutrition, University of Hohenheim, Stuttgart, Germany. Laith J. Abu-Raddad, PhD, Infectious Disease Epidemiology Group, Weill Cornell Medical College in Qatar, Doha, Qatar. Kelemework Adane, MS, Department of Medical Microbiology and Immunology, College of Health Sciences, Mekelle University, Mekelle, Ethiopia. Aliasghar Ahmad Kiadaliri, PhD, Department of Clinical Sciences Lund, Orthopedics, Clinical Epidemiology Unit, Lund University, Lund, Sweden. Alireza Ahmadi, PhD, Kermanshah University of Medical Sciences, Kermanshah, Iran. Muktar Beshir, MPH, College of Health Sciences, Department of Epidemiology, ICT and e-Learning Coordinator, Jimma University, Jimma, Ethiopia. Ayman Al-Eyadhy, MD, King Saud University, Riyadh, Saudi Arabia. Khurshid Alam, PhD, Murdoch Childrens Research Institute, The University of Melbourne, Parkville, Victoria, Australia; The University of Melbourne, Melbourne, VIC, Australia; The University of Sydney, Sydney, NSW, Australia. Noore Alam, MAppEpid, Department of Health, Queensland, Brisbane, Queensland, Australia. Deena

- Alasfoor, MSc, Ministry of Health, Al Khuwair, Oman. Reza Alizadeh-Navaei, PhD, Gastrointestinal Cancer Research Center, Mazandaran University of Medical Sciences, Sari, Iran. Fatma Al-Maskari, PhD, College of Medicine and Health Sciences, United Arab Emirates University, Al-Ain City, United Arab Emirates. Rajaa Al-Raddadi, PhD, Joint Program of Family and Community Medicine, Jeddah, Saudi Arabia. Ubai Alsharif, MPH, Charité Universitätsmedizin, Berlin, Germany. Khalid A. Altirkawi, MD, King Saud University, Riyadh, Saudi Arabia. Nahla Anber, PhD, Mansoura University, Mansoura, Egypt. Hossein Ansari, PhD, Health Promotion Research Center, Department of Epidemiology and Biostatistics, Zahedan University of Medical Sciences, Zahedan, Iran. Carl Abelardo T. Antonio, MD, Department of Health Policy and Administration, College of Public Health, University of the Philippines Manila, Manila, Philippines. Palwasha Anwari, MD, Self-employed, Kabul, Afghanistan. Hamid Asayesh, PhD, Department of Medical Emergency, School of Paramedic, Qom University of Medical Sciences, Qom, Iran. Solomon Weldegebreal Asgedom, PhD, Mekelle University, Mekelle, Ethiopia. Tesfay Mehari Atey, MS, Mekelle University, Mekelle, Ethiopia. Euripide Frinel G. Arthur, MPH, Institut de Recherche Clinique du Bénin (IRCB), Cotonou, Benin, Laboratoire d'Etudes et de Recherche-Action en Santé (LERAS Afrique), Parakou, Benin. Umar Bacha, PhD, School of Health Sciences, University of Management and Technology, Lahore, Pakistan. Aleksandra Barac, PhD, Faculty of Medicine, University of Belgrade, Belgrade, Serbia. Shahrzad Bazargan-Hejazi, PhD, College of Medicine, Charles R. Drew University of Medicine and Science, Los Angeles, CA, United States; David Geffen School of Medicine, University of California at Los Angeles, Los Angeles, CA, United States. Neeraj Bedi, MD, College of Public Health and Tropical Medicine, Jazan, Saudi Arabia. Zulfiqar A. Bhutta, PhD, Centre of Excellence in Women and Child Health, Aga Khan University, Karachi, Pakistan; Centre for Global Child Health, The Hospital for Sick Children, Toronto, ON, Canada. Michael Brauer, ScD, University of British Columbia, Vancouver, BC, Canada; Institute for Health Metrics and Evaluation, University of Washington, Seattle, Washington, United States. Zahid A Butt, PhD, Al Shifa Trust Eye Hospital, Rawalpindi, Pakistan. Carlos A. Castañeda-Orjuela, MSc, Colombian National Health Observatory, Instituto Nacional de Salud, Bogota, Colombia; Epidemiology and Public Health Evaluation Group, Public Health Department, Universidad Nacional de Colombia, Bogota, Colombia. Hadi Danawi, PhD, Walden University, Minneapolis, Minnesota, United States. Shirin Djalalinia, PhD, Undersecretary for Research and Technology, Ministry of Health and Medical Education, Tehran, Iran. Aman Yesuf Endries, MPH, Arba Minch University, Arba Minch, Ethiopia. Babak Eshрати, PhD, Ministry of Health and Medical Education, Tehran, Iran, Arak University of Medical Sciences, Arak, Iran. Maryam S Farvid, PhD, Department of Nutrition, Harvard T. H. Chan School of Public Health, Harvard University, Boston, MA, United States; Harvard/MGH Center on Genomics, Vulnerable Populations, and Health Disparities, Mongan Institute for Health Policy, Massachusetts General Hospital, Boston, MA, United States. Seyed-Mohammad Fereshtehnejad, PhD, Department of Neurobiology, Care Sciences and Society (NVS), Karolinska Institutet, Stockholm, Sweden. Florian Fischer, PhD, School of Public Health, Bielefeld University, Bielefeld, Germany; Alberto L. Garcia-Basteiro, MSc Epidemiology, Manhica Health Research Center, Manhica, Mozambique, Barcelona Institute for Global Health, Barcelona, Spain. Kiros Tedla Gebrehiwot, MSc, Mekelle University, Mekelle, Ethiopia. Tsegaye Tewelde Gebrehiwot, MPH, Jimma University, Jimma, Ethiopia. Gessesew Bugssa Hailu, MSc, Mekelle University, Mekelle, Ethiopia, Kilte Awlaelo Health and Demographic Surveillance System, Mekelle, Ethiopia. Randah Ribhi Hamadeh, DPhil, Arabian Gulf University, Manama, Bahrain. Mitiku Teshome Hambisa, MPH, College of Health and Medical Sciences, Haramaya University, Harar, Ethiopia. Samer Hamidi, DrPH, Hamdan Bin Mohammed Smart University, Dubai, United Arab Emirates. Mohammad Sadegh Hassanvand, PhD, Center for Air Pollution Research, Institute for Environmental Research, Tehran University of Medical Sciences, Tehran, Iran. Mohammad T. Hedayati, PhD, Department of Medical Mycology and Parasitology, School of Medicine, Mazandaran University of Medical Sciences, Sari, Iran. Nobuyuki Horita, MD, Department of Pulmonology, Yokohama City University Graduate School of Medicine, Yokohama, Japan. Abdullatif Husseini, PhD, Institute of Community and Public Health, Birzeit University, Birzeit, Palestine. Spencer Lewis James, MD, Denver Health/University of Colorado, Denver, CO, United States. Mehdi Javanbakht PhD, University of Aberdeen, Aberdeen, United Kingdom. Jost B. Jonas, MD, Department of Ophthalmology, Medical Faculty Mannheim, Ruprecht-Karls-University Heidelberg, Mannheim, Germany. Amir Kasaeian, PhD, Hematology-Oncology and Stem Cell Transplantation Research Center, Tehran University of Medical Sciences, Tehran, Iran; Endocrinology and Metabolism Population Sciences Institute, Tehran University of Medical Sciences, Tehran, Iran. Yousef Saleh Khader, ScD, Department of Community Medicine, Public Health and Family Medicine, Jordan University of Science and Technology, Irbid, Jordan. Ejaz Ahmad Khan, MD, Health Services Academy, Islamabad, Pakistan. Gulfaraz Khan, PhD, Department of Microbiology and Immunology, College of Medicine and Health Sciences, United Arab Emirates University, Al Ain, United Arab Emirates. Abdullah Khoja, MD, Mohammed Ibn Saudi University, Riyadh, Saudi Arabia. Jagdish Khubchandani, PhD, Department of Nutrition and Health Science, Ball State University, Muncie, Indiana, United States. Yun Jin Kim, PhD, Faculty of Chinese Medicine, Southern University College, Skudai, Malaysia. Niranjana Kissoon, MD, University of British Columbia, Vancouver, BC, Canada. Heidi J. Larson, PhD, Department of Infectious Disease Epidemiology, London School of Hygiene and Tropical Medicine, London, United Kingdom; Institute for Health Metrics and Evaluation, University of Washington, Seattle, Washington, United States. Asma Abdul Latif, PhD, Department of Zoology, Lahore College for Women University, Lahore, Pakistan. Cheru Tesema Leshargie, MPH, Debre Markos University, Debre Markos, Ethiopia. Raimundas Lunevicius, PhD, Aintree University Hospital National Health Service Foundation Trust, Liverpool, United Kingdom; School of Medicine, University of Liverpool, Liverpool, United Kingdom. Hassan Magdy Abd El Razek, MBBCH, Mansoura Faculty of Medicine, Mansoura, Egypt. Mohammed Magdy Abd El Razek, MBBCH, Aswan University Hospital, Aswan Faculty of Medicine, Aswan, Egypt. Reza Majdzadeh, PhD, Knowledge Utilization Research Center and Community Based Participatory Research Center, Tehran University of Medical Sciences, Tehran, Iran. Azeem Majeed, MD, Department of Primary Care and Public Health, Imperial College London, London, United Kingdom. Reza Malekzadeh, MD, Digestive Diseases Research Institute, Tehran University of Medical Sciences, Tehran, Iran; Digestive Diseases Research Institute, Tehran University of Medical Sciences, Tehran, Iran. Habibolah Masoudi Farid, MD, State Welfare Organisation, Tehran, Iran. Alem Mehari, MD, College of Medicine, Howard University, Washington, DC, United States. Ziad A. Memish, MD, Saudi Ministry of Health, Riyadh, Saudi Arabia, College of Medicine, Alfaisal University, Riyadh, Saudi Arabia. Desalegn Tadesse Mengistu, MS, College of Health Sciences, Mekelle University, Mekelle, Ethiopia. Haftay Berhane Mezgebe, MS, Mekelle University, Mekelle, Ethiopia. Sachiko Nakamura, MSc, Theoretical Biology and Medical Modelling, Graduate School of Medicine, Hokkaido University, Hokkaido, Japan. Eyal Oren PhD, University of Arizona, Tucson, AZ, United States. Mahesh PA, DNB, JSS Medical College, JSS University, Mysore, India. Farshad Pourmalek, PhD, University of British Columbia, Vancouver, British Columbia, Canada. Mostafa Qorbani, PhD, Non-communicable Diseases Research Center, Alborz University of Medical Sciences, Karaj, Iran. Amir Radfar, MD, A T Still University, Kirksville, MO, United

States. Anwar Rafay, MS, Contech International Health Consultants, Lahore, Pakistan; Contech School of Public Health, Lahore, Pakistan. Vafa, Rahimi-Movaghar, MD, Sina Trauma and Surgery Research Center, Tehran University of Medical Sciences, Tehran, Iran. Rajesh Kumar Rai, MPH, Society for Health and Demographic Surveillance, Suri, India. David Laith Rawaf, MD, WHO Collaborating Centre, Imperial College London, London, United Kingdom; North Hampshire Hospitals, Basingstoke, United Kingdom; University College London Hospitals, London, United Kingdom. Salman Rawaf MD, Imperial College London, London, United Kingdom. Amany H Refaat, PhD, Walden University, Minneapolis, MN, United States; Suez Canal University, Ismailia, Egypt. Satar Rezaei, PhD, School of Public Health, Kermanshah University of Medical Sciences, Kermanshah, Iran. Mohammad Sadegh Rezaei, MD, Infectious Diseases Research Centre with Focus on Nosocomial Infection, Mazandaran University of Medical Sciences, Sari, Iran. Hirbo Shore Roba, MPH, College of Health and Medical Sciences, Haramaya University, Harar, Ethiopia. Gholamreza Roshandel, PhD, Golestan Research Center of Gastroenterology and Hepatology, Golestan University of Medical Sciences, Gorgan, Iran; Digestive Diseases Research Institute, Tehran University of Medical Sciences, Tehran, Iran. Mahdi Safdarian, MD, Sina Trauma and Surgery Research Center, Tehran University of Medical Sciences, Tehran, Iran. Saeid Safiri, PhD, Managerial Epidemiology Research Center, Department of Public Health, School of Nursing and Midwifery, Maragheh University of Medical Sciences, Maragheh, Iran. Mohammad Ali Sahraian, MD, MS Research Center, Neuroscience Institute, Tehran University of Medical Sciences, Tehran, Iran. Payman Salamati, MD, Sina Trauma and Surgery Research Center, Tehran University of Medical Sciences, Tehran, Iran. Abdallah M Samy, PhD, Ain Shams University, Cairo, Egypt. Benn Sartorius, PhD, Public Health Medicine, School of Nursing and Public Health, University of KwaZulu-Natal, Durban, South Africa; UKZN Gastrointestinal Cancer Research Centre, South African Medical Research Council (SAMRC), Durban, South Africa. Sadaf G. Sepanlou, PhD, Digestive Diseases Research Institute, Tehran University of Medical Sciences, Tehran, Iran. Masood Ali Shaikh, MD, Independent Consultant, Karachi, Pakistan. Morteza Shamsizadeh, MPH, Department of Medical Surgical Nursing, School of Nursing and Midwifery, Hamadan University of Medical Sciences, Hamadan, Iran. Mika Shigematsu, PhD, National Institute of Infectious Diseases, Tokyo, Japan, Sandia National Laboratories, Albuquerque, New Mexico, United States. Jasvinder A. Singh, MD, University of Alabama at Birmingham and Birmingham Veterans Affairs Medical Center, Birmingham, Alabama, United States. Muawiyah Babale Sufiyan, MBA, Ahmadu Bello University, Zaria, Zaria, Nigeria. Arash Tehrani-Banihashemi, PhD, Preventive Medicine and Public Health Research Center, Iran University of Medical Sciences, Tehran, Iran. Mohamad-Hani Temsah, MD, King Saud University, Riyadh, Saudi Arabia. Roman Topor-Madry, PhD, Institute of Public Health, Faculty of Health Sciences, Jagiellonian University Medical College, Kraków, Poland; Faculty of Health Sciences, Wrocław Medical University, Wrocław, Poland. Olalekan A Uthman, PhD, Warwick Medical School, University of Warwick, Coventry, United Kingdom. Stein Emil Vollset, DrPH, Center for Disease Burden, Norwegian Institute of Public Health, Bergen, Norway; Department of Global Public Health and Primary Care, University of Bergen, Bergen, Norway; Institute for Health Metrics and Evaluation, University of Washington, Seattle, Washington, United States. Fiseha Wadilo, MS, Wolaita Sodo University, Wolaita Sodo, Ethiopia. Tolassa Wakayo, MS, Jimma University, Jimma, Ethiopia. Andrea Werdecker, PhD, Competence Center Mortality-Follow-Up of the German National Cohort, Federal Institute for Population Research, Wiesbaden, Germany. Tissa Wijeratne, MD, Western Health, Footscray, Victoria, Australia,

University of Melbourne, Footscray, Victoria, Australia. Mohsen Yaghoubi, MSc, School of Public Health, University of Saskatchewan, Saskatoon, Saskatchewan, Canada. Hassen Hamid Yimam, MPH, Mizan Tepi University, Mizan Teferi, Ethiopia. Naohiro Yonemoto, MPH, Department of Biostatistics, School of Public Health, Kyoto University, Kyoto, Japan. Mustafa Z Younis, DrPH, Jackson State University, Jackson, MS, United States. Maysaa El Sayed Zaki, PhD, Faculty of Medicine, Mansoura University, Mansoura, Egypt. Aisha O. Jumaan, PhD Independent Consultant, Seattle, Washington, United States. Theo Vos, PhD, Institute for Health Metrics and Evaluation, University of Washington, Seattle, Washington, United States. Simon I. Hay, DSc, Oxford Big Data Institute, Li Ka Shing Centre for Health Information and Discovery, University of Oxford, Oxford United Kingdom; Institute for Health Metrics and Evaluation, University of Washington, Seattle, Washington, United States. Mohsen Naghavi, PhD, Institute for Health Metrics and Evaluation, University of Washington, Seattle, Washington, United States. Christopher J. L. Murray, DPhil, Institute for Health Metrics and Evaluation, University of Washington, Seattle, Washington, United States. Ali H. Mokdad, PhD, Institute for Health Metrics and Evaluation, University of Washington, Seattle, Washington, United States.

The authors would like to acknowledge Adrienne Chew at the Institute for Health Metrics and Evaluation, Seattle, WA, for editing this paper. This research was funded by the Bill and Melinda Gates Foundation.

Compliance with ethical standards

This manuscript reflects original work that has not previously been published in whole or in part and is not under consideration elsewhere. All authors have read the manuscript and have agreed that the work is ready for submission and accept responsibility for its contents. The authors of this paper have complied with all ethical standards and do not have any conflicts of interest to disclose at the time of submission. The funding source played no role in the design of the study, the analysis and interpretation of data, and the writing of the paper.

Conflict of interest Dr. Carl Abelardo T. Antonio reports Grants and personal fees from Johnson and Johnson (Philippines), Inc. outside the submitted work. Dr. Jasvinder Singh serves as the principal investigator for an investigator-initiated study funded by Horizon Pharmaceuticals through a grant to DINORA, Inc., a 501 (c)(3) entity. Dr. Singh is a member of the executive of OMERACT, an organization that develops outcome measures in rheumatology and receives arms-length funding from 36 companies; a member of the American College of Rheumatology's (ACR) Annual Meeting Planning Committee (AMPC); Chair of the ACR Meet-the-Professor, Workshop and Study Group Subcommittee; and a member of the Veterans Affairs Rheumatology Field Advisory Committee. Other authors have no conflict of interest.

Human and animal rights statement The study did not involve human participants and/or animals.

Informed consent No informed consent was needed.

Open Access This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

References

- Akseer N, Kamali M, Husain S et al (2015) Strategies to avert preventable mortality among mothers and children in the Eastern Mediterranean Region: new initiatives, new hope. *East Mediterr Health J* 21:361–373
- Assefa F, Jabarkhil MZ, Salama P, Spiegel P (2001) Malnutrition and mortality in Kohistan district, Afghanistan, April 2001. *J Am Med Assoc* 286:2723–2728
- Beauté J, Snacken R, Adlhoch C (2014) Annual epidemiological report, respiratory tract infections. <https://ecdc.europa.eu/sites/portal/files/media/en/publications/Publications/Respiratory-tract-infections-annual-epidemiological-report-2014.pdf>. Accessed 11 July 2017
- Bhutta ZA, Das JK, Walker N et al (2013) Interventions to address deaths from childhood pneumonia and diarrhoea equitably: what works and at what cost? *Lancet Lond Engl* 381:1417–1429. doi:10.1016/S0140-6736(13)60648-0
- Breiman RF, Van Beneden CA, Farnon EC (2013) Surveillance for respiratory infections in low- and middle-income countries: experience from the centers for disease control and prevention's global disease detection international emerging infections program. *J Infect Dis* 208:S167–S172. doi:10.1093/infdis/jit462
- Burki T (2015) Yemen health situation “moving from a crisis to a disaster”. *Lancet Lond Engl* 385:1609. doi:10.1016/S0140-6736(15)60779-6
- Burki T (2016) Yemen's neglected health and humanitarian crisis. *Lancet* 387:734–735. doi:10.1016/S0140-6736(16)00389-5
- Cohen AJ, Brauer M, Burnett R et al (2017) Estimates and 25-year trends of the global burden of disease attributable to ambient air pollution: an analysis of data from the Global Burden of Diseases Study 2015. *Lancet Lond Engl* 389:1907–1918. doi:10.1016/S0140-6736(17)30505-6
- Dewachi O, Skelton M, Nguyen V-K et al (2014) Changing therapeutic geographies of the Iraqi and Syrian wars. *Lancet Lond Engl* 383:449–457. doi:10.1016/S0140-6736(13)62299-0
- Forouzanfar MH, Afshin A, Alexander LT et al (2016) Global, regional, and national comparative risk assessment of 79 behavioural, environmental and occupational, and metabolic risks or clusters of risks, 1990–2015: a systematic analysis for the Global Burden of Disease Study 2015. *Lancet* 388:1659–1724. doi:10.1016/S0140-6736(16)31679-8
- Framework Convention Alliance Parties to the WHO FCTC (2017). <http://www.fctc.org/about-fca/tobacco-control-treaty/latest-ratifications/parties-ratifications-accessions>. Accessed 6 Jun 2017
- GAVI (2017) The vaccine alliance Pneumococcal vaccine support. <http://www.gavi.org/support/nvs/pneumococcal/>. Accessed 11 July 2017
- Guha-Sapir D, Panhuis WG (2004) Conflict-related mortality: an analysis of 37 datasets. *Disasters* 28:418–428. doi:10.1111/j.0361-3666.2004.00267.x
- Hadi A (2003) Management of acute respiratory infections by community health volunteers: experience of Bangladesh Rural Advancement Committee (BRAC). *Bull World Health Organ* 81:183–189
- Haq Z, Mahjour J, Khan W (2013) Communicable diseases in the Eastern Mediterranean Region: prevention and control 2010–2011. *East Mediterr Health J* 19:888–891
- Global Burden of Disease Study (GBD) (2015) Institute for Health Metrics and Evaluation. Data Input Source Tool. <http://ghdx.healthdata.org/gbd-2015/data-input-sources>
- Kassebaum NJ, Arora M, Barber RM et al (2016) Global, regional, and national disability-adjusted life-years (DALYs) for 315 diseases and injuries and healthy life expectancy (HALE), 1990–2015: a systematic analysis for the Global Burden of Disease Study 2015. *Lancet* 388:1603–1658. doi:10.1016/S0140-6736(16)31460-X
- Kheirandish M, Rashidian A, Kebriaeezade A et al (2015) A review of pharmaceutical policies in response to economic crises and sanctions. *J Res Pharm Pract* 4:115–122. doi:10.4103/2279-042X.162361
- Kovacs SD, Mullholland K, Bosch J et al (2015) Deconstructing the differences: a comparison of GBD 2010 and CHERG's approach to estimating the mortality burden of diarrhea, pneumonia, and their etiologies. *BMC Infect Dis* 15:16. doi:10.1186/s12879-014-0728-4
- Kronfol NM (2012) Access and barriers to health care delivery in Arab countries: a review. *East Mediterr Health J* 18:1239–1246
- Levy BS, Sidel VW (2016) Documenting the effects of armed conflict on population health. *Annu Rev Public Health* 37:205–218. doi:10.1146/annurev-publhealth-032315-021913
- Massoumi RL, Koduri S (2015) Adverse effects of political sanctions on the health care system in Iran. *J Glob Health* 5:020302. doi:10.7189/jogh.05-020302
- Milne-Price S, Miazgowiec KL, Munster VJ (2014) The emergence of the Middle East respiratory syndrome coronavirus. *Pathog Dis* 71:121–136. doi:10.1111/2049-632X.12166
- Mokdad AH, Jaber S, Aziz MI et al (2014) The state of health in the Arab world, 1990–2010: an analysis of the burden of diseases, injuries, and risk factors. *Lancet* 383(9914):309–320
- Mokdad AH, Forouzanfar MH, Daoud F et al (2016) Health in times of uncertainty in the eastern Mediterranean region, 1990–2013: a systematic analysis for the Global Burden of Disease Study 2013. *Lancet Glob Health* 4(10):e704–e713
- Moradi-Lakeh M, Esteghamati A (2013) National immunization program in Iran. *Hum Vaccines Immunother* 9:112–114. doi:10.4161/hv.22521
- Moradi-Lakeh M, El Bcheraoui C, Tuffaha M et al (2015) Tobacco consumption in the Kingdom of Saudi Arabia, 2013: findings from a national survey. *BMC Public Health* 15:611. doi:10.1186/s12889-015-1902-3
- Murray C, King G, Lopez A et al (2008) Armed Conflict as a Public Health Problem. Social Science Research Network, Rochester
- Nair H, Simões EA, Rudan I et al (2013) Global and regional burden of hospital admissions for severe acute lower respiratory infections in young children in 2010: a systematic analysis. *Lancet* 381:1380–1390. doi:10.1016/S0140-6736(12)61901-1
- Nichol KL, Nordin JD, Nelson DB et al (2007) Effectiveness of influenza vaccine in the community-dwelling elderly. *N Engl J Med* 357:1373–1381. doi:10.1056/NEJMoa070844
- Oberg M, Jaakkola MS, Woodward A et al (2011) Worldwide burden of disease from exposure to second-hand smoke: a retrospective analysis of data from 192 countries. *Lancet Lond Engl* 377:139–146. doi:10.1016/S0140-6736(10)61388-8
- Qazi S, Aboubaker S, MacLean R et al (2015) Ending preventable child deaths from pneumonia and diarrhoea by 2025. Development of the integrated global action plan for the prevention and control of pneumonia and diarrhoea. *Arch Dis Child* 100:S23–S28. doi:10.1136/archdischild-2013-305429
- Rashidian A (2008) Adapting valid clinical guidelines for use in primary care in low and middle income countries. *Prim Care Respir J* 17:136–137. doi:10.3132/pcrj.2008.00055
- Rehfuess EA, Tzala L, Best N et al (2009) Solid fuel use and cooking practices as a major risk factor for ALRI mortality among African children. *J Epidemiol Community Health* 63:887–892. doi:10.1136/jech.2008.082685
- Renzetti G, Silvestre G, D'Amaro C et al (2009) Less air pollution leads to rapid reduction of airway inflammation and improved airway function in asthmatic children. *Pediatrics* 123:1051–1058. doi:10.1542/peds.2008-1153
- Roth DE, Caulfield LE, Ezzati M, Black RE (2008) Acute lower respiratory infections in childhood: opportunities for reducing

- the global burden through nutritional interventions. *Bull World Health Organ* 86:356–364. doi:[10.2471/BLT.07.049114](https://doi.org/10.2471/BLT.07.049114)
- Roth DE, Richard SA, Black RE (2010) Zinc supplementation for the prevention of acute lower respiratory infection in children in developing countries: meta-analysis and meta-regression of randomized trials. *Int J Epidemiol* 39:795–808. doi:[10.1093/ije/dyp391](https://doi.org/10.1093/ije/dyp391)
- Rudan I, Boschi-Pinto C, Biloglav Z et al (2008) Epidemiology and etiology of childhood pneumonia. *Bull World Health Organ* 86:408–416
- Rudan I, O'Brien KL, Nair H et al (2013) Epidemiology and etiology of childhood pneumonia in 2010: estimates of incidence, severe morbidity, mortality, underlying risk factors and causative pathogens for 192 countries. *J Glob Health* 3:010401. doi:[10.7189/jogh.03.010401](https://doi.org/10.7189/jogh.03.010401)
- Shah UH, Abu-Shaheen AK, Malik MA et al (2013) The efficacy of zinc supplementation in young children with acute lower respiratory infections: a randomized double-blind controlled trial. *Clin Nutr Edinb Scotl* 32:193–199. doi:[10.1016/j.clnu.2012.08.018](https://doi.org/10.1016/j.clnu.2012.08.018)
- Sibak M, Moussa I, El-Tantawy N et al (2015) Cost-effectiveness analysis of the introduction of the pneumococcal conjugate vaccine (PCV-13) in the Egyptian national immunization program, 2013. *Vaccine* 33(Suppl 1):A182–A191. doi:[10.1016/j.vaccine.2014.12.044](https://doi.org/10.1016/j.vaccine.2014.12.044)
- Takian A, Doshmangir L, Rashidian A (2013) Implementing family physician programme in rural Iran: exploring the role of an existing primary health care network. *Fam Pract* 30:551–559. doi:[10.1093/fampra/cmt025](https://doi.org/10.1093/fampra/cmt025)
- Tong N (2013) Background Paper 6.22—Pneumonia. http://www.who.int/medicines/areas/priority_medicines/BP6_22Pneumo.pdf. Accessed 11 July 2017
- Usmanova G, Mokdad AH (2013) Results of the global youth tobacco survey and implementation of the WHO framework convention on tobacco control in the WHO Eastern Mediterranean Region (EMR) countries. *J Epidemiol Glob Health* 3:217–234. doi:[10.1016/j.jegh.2013.07.001](https://doi.org/10.1016/j.jegh.2013.07.001)
- Voordouw BCG, van der Linden PD, Simonian S et al (2003) Influenza vaccination in community-dwelling elderly: impact on mortality and influenza-associated morbidity. *Arch Intern Med* 163:1089–1094. doi:[10.1001/archinte.163.9.1089](https://doi.org/10.1001/archinte.163.9.1089)
- Vos T, Allen C, Arora M et al (2016) Global, regional, and national incidence, prevalence, and years lived with disability for 310 diseases and injuries, 1990–2015: a systematic analysis for the Global Burden of Disease Study 2015. *Lancet* 388:1545–1602. doi:[10.1016/S0140-6736\(16\)31678-6](https://doi.org/10.1016/S0140-6736(16)31678-6)
- Wang H, Naghavi M, Allen C et al (2016) Global, regional, and national life expectancy, all-cause mortality, and cause-specific mortality for 249 causes of death, 1980–2015: a systematic analysis for the Global Burden of Disease Study 2015. *Lancet* 388:1459–1544. doi:[10.1016/S0140-6736\(16\)31012-1](https://doi.org/10.1016/S0140-6736(16)31012-1)
- WHO Regional Office for the Eastern Mediterranean (2016) The work of WHO in the Eastern Mediterranean Region: annual report of the regional director 2015. <http://www.emro.who.int/annual-report/2015/index.html>. Accessed 11 July 2017
- Williams DJ, Shah SS (2012) Community-acquired pneumonia in the conjugate vaccine era. *J Pediatr Infect Dis Soc* 1:314–328. doi:[10.1093/jpids/pis101](https://doi.org/10.1093/jpids/pis101)
- Williams N, Woodward H, Majeed A, Saxena S (2011) Primary care strategies to improve childhood immunisation uptake in developed countries: systematic review. *JRSM Short Rep* 2:81. doi:[10.1258/shorts.2011.011112](https://doi.org/10.1258/shorts.2011.011112)
- World Health Organization (2012) Vaccines against influenza WHO position paper. http://www.who.int/immunization/position_papers/PP_influenza_november2012_summary.pdf. Accessed 11 July 2017
- World Health Organization (2017) Haemophilus influenzae type b third dose (Hib3) immunization coverage. <http://www.who.int/gho/immunization/Hib/en/>. Accessed 11 July 2017
- World Health Organization Eastern Mediterranean Region (2017) Addressing the impact of air pollution on health in the Eastern Mediterranean Region. <http://www.emro.who.int/about-who/rc61/impact-air-pollution.html>. Accessed 6 June 2017
- World Health Organization and UNICEF (1986) Basic principles for control of acute respiratory infections in children in developing countries. http://apps.who.int/iris/bitstream/10665/39683/1/9241560940_eng.pdf. Accessed 11 July 2017
- World Health Organization and UNICEF (2013) Ending preventable child deaths from pneumonia and diarrhoea by 2025: The integrated global action plan for pneumonia and diarrhoea. http://www.who.int/maternal_child_adolescent/documents/global_action_plan_pneumonia_diarrhoea/en/. Accessed 11 July 2017

Electronic Supplementary Material

Article title:

Burden of lower respiratory infections in the Eastern Mediterranean Region between 1990 and 2015: Findings from the Global Burden of Disease 2015 study

Journal:

International Journal of Public Health

Authors:

GBD 2015 Eastern Mediterranean Region LRI Collaborators

Corresponding author:

Ali H. Mokdad

Institute for Health Metrics and Evaluation, University of Washington, Seattle, WA, United States

Email: mokdaa@uw.edu

e-Table 1. Age-standardized sex-specific incidence rates of lower respiratory infections (percent) in the countries of the Eastern Mediterranean Region, 2015. (Global Burden of Disease Study 2015, Eastern Mediterranean Countries, 2015)

Countries	Male			Female			Both		
	Incidence	95% UI		Incidence	95% UI		Incidence	95% UI	
Afghanistan	4.0	4.3	3.8	6.0	6.5	5.6	5.0	5.4	4.7
Bahrain	3.8	4.0	3.5	4.4	4.7	4.2	4.1	4.3	3.8
Djibouti	7.1	7.6	6.6	8.4	9.0	7.8	7.8	8.3	7.3
Egypt	4.2	4.5	4.0	4.7	5.0	4.4	4.5	4.7	4.2
Iran	4.1	4.4	3.9	4.8	5.1	4.5	4.4	4.7	4.2
Iraq	4.8	5.1	4.5	5.5	5.9	5.1	5.2	5.5	4.8
Jordan	4.1	4.4	3.9	4.6	4.8	4.3	4.3	4.6	4.1
Kuwait	4.0	4.2	3.8	5.0	5.3	4.7	4.4	4.7	4.2
Lebanon	4.2	4.5	3.9	5.0	5.4	4.7	4.6	4.9	4.3
Libya	4.3	4.6	4.0	5.5	5.8	5.1	4.9	5.2	4.6
Morocco	5.1	5.4	4.8	5.5	5.9	5.2	5.3	5.6	5.0
Oman	4.0	4.3	3.8	5.1	5.5	4.8	4.5	4.7	4.2
Pakistan	5.6	6.0	5.3	5.2	5.6	4.9	5.4	5.7	5.1
Palestine	4.9	5.2	4.6	5.4	5.8	5.0	5.1	5.4	4.8
Qatar	3.6	3.8	3.3	4.4	4.7	4.1	3.9	4.1	3.6
Saudi Arabia	3.6	3.8	3.5	4.4	4.6	4.2	4.0	4.2	3.8
Somalia	7.4	7.9	6.9	8.3	8.8	7.8	7.9	8.3	7.4
Sudan	5.9	6.4	5.5	6.9	7.5	6.4	6.4	6.9	6.0
Syria	4.6	5.0	4.3	5.4	5.8	5.0	5.0	5.4	4.7
Tunisia	3.1	3.3	2.9	3.7	3.9	3.4	3.4	3.6	3.2
United Arab Emirates	2.8	2.9	2.6	3.4	3.6	3.2	3.0	3.2	2.8
Yemen	7.7	8.2	7.2	8.7	9.4	8.2	8.2	8.8	7.7

Title: Burden of lower respiratory infections in the Eastern Mediterranean Region between 1990 and 2015: findings from the Global Burden of Disease 2015 study
Group title: GBD 2015 Eastern Mediterranean Region LRI Collaborators

Prefix	First Name:	Middle Na	Middle Na	Last Name:	Email	Address	Department	Institution/Organization	Institution	Institution	Institution	Department 2	Institution/Organization 2	Institution	Institution/O	Institution	Department	Institution	Institution	Institution	
Dr.	Maziar			Moradi-Lakeh	mazmorad	MD	Department of Community Medicine, Gastrointestinal	Iran University of Medical Sciences	Tehran												
Dr.	Charbel			El Bcheraoui	charbel@u	PhD	Institute for Health Metrics and Evaluation	University of Washington	Seattle	Washington	United_States										
Dr.	Raghd			Charara	raghd@u	MD		American University of Beirut	Beirut												
Dr.	Ibrahim			Khalil	ikhallil@uw	MD	Institute for Health Metrics and Evaluation	University of Washington	Seattle	Washington	United_States										
Dr.	Ashkan			Afshin	ashkan.afs	MD	Institute for Health Metrics and Evaluation	University of Washington	Seattle	Washington	United_States										
Dr.	Nicholas	J		Kassebaum	nickjk@uw	MD	Institute for Health Metrics and Evaluation	University of Washington	Seattle	Washington	United_States	Department of Anesthesiology & Pain Medicine	Seattle Children's Hospital	Seattle	Washington	United_States					
Mr.	Michael			Collison	m1c314@u	BS	Institute for Health Metrics and Evaluation	University of Washington	Seattle	Washington	United_States										
Ms.	Farah			Daoud	farahd@u	BA/BS	Institute for Health Metrics and Evaluation	University of Washington	Seattle	Washington	United_States										
Ms.	Adrienne			Chew	apchew@u	ND	Institute for Health Metrics and Evaluation	University of Washington	Seattle	Washington	United_States										
Mr.	Kristopher	J.		Krohn	kkrohn1@	BA	Institute for Health Metrics and Evaluation	University of Washington	Seattle	Washington	United_States										
Dr.	Danny			Colombara	dvc2@uw.	PhD	Institute for Health Metrics and Evaluation	University of Washington	Seattle	Washington	United_States										
Ms.	Rebecca			Ehrenkranz	rehrenk1@	MPH	Institute for Health Metrics and Evaluation	University of Washington	Seattle	Washington	United_States										
Dr.	Kyle	J.		Foreman	kfor@uw.	PhD	Institute for Health Metrics and Evaluation	University of Washington	Seattle	Washington	United_States		Imperial College London	London							United_Kingdom
Mr.	Joseph			Frostad	jfrostad@	MPH	Institute for Health Metrics and Evaluation	University of Washington	Seattle	Washington	United_States										
Mr.	William	W.		Godwin	wgodwin@	BS	Institute for Health Metrics and Evaluation	University of Washington	Seattle	Washington	United_States										
Mr.	Michael			Kutz	kutz13@u	BS	Institute for Health Metrics and Evaluation	University of Washington	Seattle	Washington	United_States										
Ms.	Puja	C		Rao	pujara@u	MPH	Institute for Health Metrics and Evaluation	University of Washington	Seattle	Washington	United_States										
Dr.	Robert			Reiner	rbreiner@	BA	Institute for Health Metrics and Evaluation	University of Washington	Seattle	Washington	United_States										
Mr.	Christopher			Troeger	ctroeger@	MPH	Institute for Health Metrics and Evaluation	University of Washington	Seattle	Washington	United_States										
Dr.	Haidong			Wang	haidong@	PhD	Institute for Health Metrics and Evaluation	University of Washington	Seattle	Washington	United_States										
Mr.	Amanuel	Alemu		Abajobir	amanuel.al	MPH	School of Public Health	University of Queensland	Brisbane	QLD	Australia										
Prof.	Kaja	M		Abbas	kaja.abbas	PhD		Virginia Tech	Blacksburg	VA	United_States										
Mr.	Semaw	Ferede		Abera	semawfer@	MSc	School of Public Health, College of Health Sciences	Mekelle University	Mekelle		Ethiopia	Food Security and Institute for Biological Chemistry and Nutrition	University of Hohenheim	Stuttgart							Germany
Dr.	Laith	J.		Abu-Raddad	lja2002@q	PhD	Infectious Disease Epidemiology Group	Weill Cornell Medical College in Qatar	Doha		Qatar										
Mr.	Kelemework			Adane	ingoldmit@	MS	Department of Medical Microbiology and Immunology, College of Health Sciences	Mekelle University	Mekelle		Ethiopia										
Dr.	Aliasghar			Ahmad Kiadaliri	aliasghar.a	PhD	Department of Clinical Sciences Lund, Orthopedics, Clinical Epidemiology Unit	Lund University	Lund		Sweden										
Dr.	Alireza			Ahmadi	ahmadiar1	PhD		Kermanshah University of Medical Sciences	Kermanshah		Iran			Stockholm							Sweden
Mr.	Muktar	Beshir		Ahmed	muktar27@	MPH	College of Health Sciences, Department of Epidemiology, ICT and e-Learning Coordinator	Jimma University	Jimma	Oromiya	Ethiopia										
Dr.	Ayman			Al- Eyadhy	aleyadhy@	MD		King Saud University	Riyadh		Saudi_Arabia										Saudi_Arabia
Dr.	Khurshid			Alam	khurshid.b	PhD	Murdoch Childrens Research Institute	The University of Melbourne	Parkville	Victoria	Australia		The University of Melbourne	Melbourne	VIC	Australia					The Univer
Mr.	Noore			Alam	noore.alan	MAppEpid		Department of Health, Queensland	Brisbane		Australia										NSW
Ms.	Deena			Alasfoor	deena.alas	MSc		Ministry of Health	Al Khuwair		Muscat	Oman									Australia
Dr.	Reza			Alizadeh-Navaei	reza_nava	PhD	Gastrointestinal Cancer Research Center	Mazandaran University of Medical Sciences	Sari		Mazandarā	Iran									
Prof.	Fatma			Al-Maskari	fatma.am@	PhD	College of Medicine & Health Sciences	United Arab Emirates University	Al-Ain City		United_arab_emirates										
Dr.	Rajaa			Al-Raddadi	saudiresea	PhD		Joint Program of Family and Community Medicine	Jeddah	Makkah	Saudi_Arabia										
Dr.	Ubai			Alsharif	ubai.al-sha	MPH		Charité Universitätsmedizin	Berlin	Berlin	Germany										
Dr.	Khalid	A.		Altrikawi	kaltirkawi@	MD		King Saud University	Riyadh		Saudi_Arabia										
Dr.	Nahla			Anber	nahlaanbe	PhD		Mansoura University	Mansoura		Egypt										
Dr.	Hossein			Ansari	haepidemi	PhD	Health Promotion Research Center, Department of Epidemiology and Biostatistics	Zahedan University of Medical Sciences	Zahedan		Iran										
Dr.	Carl Abelardo	T		Antonio	ctantonion	MD	Department of Health Policy and Administration,	University of the Philippines Manila	Manila		Philippines										
Dr.	Palwasha			Anwari	anwari222	MD	College of Public Health	Self-employed	Kabul	Kabul	Afghanistan										
Mr.	Hamid			Asayesh	asayeshps,	PhD	Department of Medical Emergency, School of Paramedic	Qom University of Medical Sciences	Qom		Iran										
Mr.	Solomon	Weldegebreal		Asgedom	solomon.a:	PhD		Mekelle University	Mekelle	Tigray	Ethiopia										
Mr.	Tesfay Mehari			Atey	tesfay.meh	MS		Mekelle University	Mekelle	Tigray	Ethiopia										
Dr.	Euripide	Frinel	G. Arthur	Avokpaho	a_euripide	MPH		Institut de Recherche Clinique du Bénin (IRCB)	Cotonou		Benin	Laboratoire d'Etudes et de Recherche-Action en Santé (LERAS Afrique)	Parakou								Benin
Mr.	Umar			Bacha	umar.bach	PhD	School of Health Sciences	University of Management and Technology	Lahore	Punjab	Pakistan										
Dr.	Aleksandra			Barac	aleksandra	PhD	Faculty of Medicine	University of Belgrade	Belgrade	Belgrade	Serbia										
Prof.	Shahzad			Bazargan-Hejazi	shahrzadb:	PhD	College of Medicine	Charles R. Drew University of Medicine and Science	Los Angeles	CA	United_States	David Geffen School of Medicine	Los Angeles	CA							United_States
Dr.	Neeraj			Bedi	drnbedi@y	MD		College of Public Health and Tropical Medicine	Jazan		Saudi_Arabia										
Dr.	Zulfiqar	A.		Bhutta	zulfiqar.bh	PhD	Centre of Excellence in Women and Child Health	Aga Khan University	Karachi		Pakistan	Centre for Global Child Health Institute for Health Metrics and Evaluation	The Hospital for Sick Children	Toronto	ON	Canada					
Prof.	Michael			Brauer	michael.br	ScD		University of British Columbia	Vancouver	BC	Canada		University of Washington	Seattle	WA	United_States					
Dr.	Zahid	A		Butt	zabutt3@y	PhD		Al Shifa Trust Eye Hospital	Rawalpindi	Punjab	Pakistan										
Dr.	Carlos	A.		Castañeda-Orjuela	ccastanedz	MSc	Colombian National Health Observatory	Instituto Nacional de Salud	Bogota	Bogota,	DC	Colombia	Epidemiology and Public Health Evaluation Group, Public Health Department	Universidad Nacional de Colombia	Bogota	Bogota, DC	Colombia				

Dr. Mr.	Hadi Shirin Aman	Yesuf	Danawi Djalalinia Endries	hdanawi@PhD shdjaliniz PhD amanyesuf MPH	Undersecretary for Research & Technology	Walden University Ministry of Health & Medical Education Arba Minch SNNPR	Minneapolis Tehran Arba Minch	United_States Iran Ethiopia				
Dr.	Babak		Eshrati	eshratib@PhD		Ministry of Health and Medical Education	Tehran Tehran Iran			Arak University of Medical Sciences	Arak	Iran
Dr.	Maryam	S	Farvid	mfarvid@PhD	Department of Nutrition, Harvard T.H. Chan School of Public Health	Harvard University	Boston MA United_States			Harvard/MGH Center on Genomics, Vulnerable Populations, and Health Disparities, Mongan Institute for Health Policy, Massachusetts General Hospital	Boston MA United_States	
Dr. Dr.	Seyed-Mohammad Florian		Fereshtehnejad Fischer	sm.feresht PhD f.fischer@PhD	Department of Neurobiology, Care Sciences and Society (NVS) School of Public Health	Karolinska Institutet Bielefeld University	Stockholm Bielefeld	Sweden North Rhin Germany				
Dr. Mr. Mr.	Alberto Kiros Tsegaye	L Tedla Tewelde	Garcia-Basteiro Gebrehiwot Gebrehiwot	alberto.gar MSc Epidemiology Kirosmerry MSc tsegaye.te MPH		Manhiça Health Research Center Mekelle University Jimma University	Manhiça Mekelle Jimma	Maputo Tigray Oromia	Mozambique Ethiopia Ethiopia	Barcelona Institute for Global Health	Barcelona Barcelona Spain	
Mr. Prof. Mr. Dr.	Gessesew Randah Mitiku Samer	Bugssa Ribhi Teshome	Hailu Hamadeh Hambisa Hamidi	bugssag@PhD randah@PhD mitikkuuf@PhD s.hamidi@DrPH		Mekelle University Arabian Gulf University Haramaya University Hamdan Bin Mohammed Smart University	Mekelle Manama Harar Dubai	Ethiopia Bahrain Ethiopia united_arab_emirates		Kilte Awlelo Health and Demographic Surveillance System	Mekelle	Ethiopia
Dr.	Mohammad Sadegh		Hassanvand	hassanvan@PhD	Center for Air Pollution Research, Institute for Environmental Research Department of Medical Mycology and Parasitology, School of Medicine	Tehran University of Medical Sciences	Tehran Iran					
Prof.	Mohammad	T.	Hedayati	hedayatim@PhD		Mazandaran University of Medical Sciences Yokohama City University Graduate School of Medicine	Sari Mazandara Iran					
Dr. Dr. Dr. Dr.	Nobuyuki Abdullatif Spencer Mehdi	Lewis	Horita Husseini James Javanbakht	horitano@ MD abdullatif PhD spencj@gn MD javanbakht PhD	Department of Pulmonology Institute of Community and Public Health	Birzeit University Denver Health/University of Colorado University of Aberdeen	Yokohama Birzeit Denver Aberdeen	Kanagawa Palestine United_States United_Kingdom				
Prof.	Jost	B	Jonas	Jost.Jonas@ MD	Department of Ophthalmology, Medical Faculty Mannheim	Ruprecht-Karls-University Heidelberg	Mannheim Germany	Germany				
Dr.	Amir		Kasaean	amir_kasa@PhD	Hematology-Oncology and Stem Cell Transplantation Research Center Department of Community Medicine, Public Health and Family Medicine	Tehran University of Medical Sciences Jordan University of Science and Technology Health Services Academy	Tehran Tehran Iran	Endocrinology and Metabolism Population Sciences Institute	Tehran University of Medical Sciences	Tehran Tehran Iran		
Prof. Dr.	Yousef Ejaz	Saleh Ahmad	Khader Khan	yskader@ScD ejaz@hsa.ri MD		Jordan University of Science and Technology Health Services Academy	Irbid Irbid Islamabad	Jordan Pakistan				
Dr. Dr. Dr. Dr. Prof.	Gulfaraz Abdullah Jagdish Yun Jin Niranjan	Tawfih Abdullah	Khan Khoja Khubchandani Kim Kissoon	g_khan@u PhD atkhaja@ MD jkhubchani PhD neurokim7 PhD nkissoon@ MD	Department of Microbiology and Immunology, College of Medicine & Health Sciences	United Arab Emirates University Mohammed Ibn Saudi University Ball State University Southern University College University of British Columbia	Al Ain Riyadh Muncie Skudai Vancouver BC	Abu Dhabi Saudi Arabia Indiana Johor Canada	United Arab Emirates Saudi Arabia United_States Malaysia Canada			
Dr. Dr. Mr.	Heidi Asma Cheru	J. Abdul Tesema	Larson Latif Leshargie	Heidi.Larsc PhD asma5latif PhD chertesem MPH	Department of Infectious Disease Epidemiology Department of Zoology	London School of Hygiene & Tropical Medicine Lahore College for Women University Debre Markos University Aintree University Hospital National Health Service Foundation Trust Mansoura Faculty of Medicine Aswan Faculty of Medicine	London Lahore Lahore Debre Mar Amhara	United_Kin Evaluation Pakistan Ethiopia	Institute for Health Metrics and University of Washington	Seattle WA United_States		
Prof. Dr. Dr.	Raimundas Hassan Mohammed		Lunevicius Magdy Abd El Razek Magdy Abd El Razek	rlunevichu PhD farreha201 MBBCh hoda_734@MBBCh		Aswan University Hospital	Liverpool Mansoura Aswan	United_Kin School of Medicine Egypt Egypt	University of Liverpool	Liverpool	United_Kingdom	
Prof. Prof.	Reza Azeem		Majdzadeh Majeed	rezamajd@PhD a.majeed@ MD	Knowledge Utilization Research Center and Community Based Participatory Research Center Department of Primary Care & Public Health	Tehran University of Medical Sciences Imperial College London	Tehran London	Iran United_Kingdom				
Prof. Dr. Dr. Dr. Prof. Mr. Mr. Mr.	Reza Habibollah Alem Ziad Desalegn Haftay	A. Tadese Berhane	Malekzadeh Masoudi Farid Mehari Memish Mengistu Mezgebe	malek@tu MD Masoudi.Farid MD alem.mehz MD zmemish@ MD desalegn4 MS haftay.berl MS	Digestive Diseases Research Institute	Tehran University of Medical Sciences State Welfare Organisation Howard University Saudi Ministry of Health Mekelle University Mekelle University	Tehran TEHRAN Washington DC Riyadh Mekelle Mekelle	Iran Iran United_States Saudi_Arat Ethiopia Ethiopia	Digestive Diseases Research Tehran University of Medical Sciences	Tehran	Iran	
Ms. Prof. Prof. Dr.	Sachiko Eyal Mahesh Farshad		Nakamura Oren PA Pourmalek	sachikonak MSc eoren@err PhD mahesh19 DNB farshad.po PhD	Theoretical Biology and Medical Modelling, Graduate School of Medicine	Hokkaido University University of Arizona JSS University University of British Columbia	Hokkaido Tucson Mysore Vancouver	Japan United_States India British Col, Canada				
Dr. Dr.	Mostafa Amir		Qorbani Radfar	mqorbani1 PhD aradfar@a MD	Non-Communicable Diseases Research Center	Alborz University of Medical Sciences A T Still University	Karaj Kirksville MO	Iran United_States				
Dr. Prof.	Anwar Vafa		Rafay Rahimi-Movaghar	rafay.aku.ri MS v_rahimi@ MD	Sina Trauma and Surgery Research Center	Contech International Health Consultants Tehran University of Medical Sciences	Lahore Tehran	Punjab Iran	Contech School of Public Health	Lahore Punjab Pakistan		

Mr. Dr. Prof.	Rajesh Kumar David Salman Amany	Laith H	Rai Rawaf Rawaf Refaat	rajesh.iips@ MPH drawaf@ic MD, MBBS WHO Collaborating Centre s.rawaf@ir MD amany.ref@ PhD	Society for Health and Demographic Surveillance Imperial College London Imperial College London Walden University	Suri London London Minneapolis MN	West Beng India United_Kingdom United_Kingdom United_States	North Hampshire Hospitals Basingstroke	United_Kingdom	University London	United_Kin	
Dr.	Satar		Rezaei	satar.rezaei PhD	School of Public Health Infectious Diseases Research Centre with Focus on Nosocomial Infection College of Health and Medical Sciences Golestan Research Center of Gastroenterology and Hepatology	Kermanshah University of Medical Sciences	Kermanshah Iran					
Dr. Mr.	Mohammad Sadegh Hirbo	Shore	Rezaei Roba	drmsrezaei@ MD hamakiya@ MPH	Mazandaran University of Medical Sciences Haramaya University	Sari Harar	Mazandara Iran Ethiopia					
Dr. Dr.	Gholamreza Mahdi		Roshandel Safdarian	roshandel_ PhD mahdisafd@ MD	Sina Trauma & Surgery Research Center Managerial Epidemiology Research Center, Department of Public Health, School of Nursing and Midwifery	Golestan University of Medical Sciences Tehran University of Medical Sciences,	Gorgan Tehran Iran	Digestive Diseases Research Institute	Tehran University of Medical Sciences	Tehran	Iran	
Dr. Dr. Prof. Dr.	Saeid Mohammad Ali Payman Abdallah	M.	Safiri Sahraian Salamati Samy	saeidsafiri(PhD sahraian13 MD psalamati@ MD asamy.ku@ PhD	MS Research Center, Neuroscience Institute Sina Trauma and Surgery Research Center	Maragheh University of Medical Sciences Tehran University of Medical Sciences Tehran University of Medical Sciences Ain Shams University	Maragheh Tehran Tehran Cairo	Iran Iran Iran Egypt			Lawrence Kansas United_States	
Prof. Dr. Dr.	Benn Sadaf Masood	G Ali	Sartorius Sepanlou Shaikh	sartorius@ PhD sepanlou@ PhD masoodali@ MD	Public Health Medicine, School of Nursing and Public Health Digestive Diseases Research Institute	University of KwaZulu-Natal Tehran University of Medical Sciences Independent Consultant	Durban Tehran Karachi	South_Afri Research Centre Iran Pakistan	UKZN Gastrointestinal Cancer Research Centre (SAMRC)	South African Medical Research Council (SAMRC)	Durban	South_Africa
Mr. Dr.	Morteza Mika		Shamsizadeh Shigematsu	mshamsizadeh@ MPH mikas@nih PhD	Department of Medical Surgical Nursing, School of Nursing and Midwifery	Hamadan University of Medical Sciences National Institute of Infectious Diseases	Hamadan Tokyo	Iran Japan		Sandia National Laboratories	Albuquerque New Mexico	United_States
Dr. Dr.	Jasvinder Muawiyah	A. Babale	Singh Sufiyan	jasvinder.n MD sufianmb@ MBA		University of Alabama at Birmingham and Birmingham Veterans Affairs Medical Center Ahmadu Bello University, Zaria	Birmingham Alabama Zaria	United_States Nigeria				
Dr.	Arash		Tehrani-Banihashemi	tehranib@ PhD	Preventive Medicine and Public Health Research Center	Iran University of Medical Sciences	Tehran Tehran Iran					
Dr.	Mohamad-Hani		Temsah	Temsah1@ MD		King Saud University	Riyadh	Saudi_Arabia	King Faisal Specialist Hospital and Research Center	Riyadh	Saudi_Arabia	
Dr. Dr.	Roman Olalekan	A	Topor-Madry Uthman	roman.top PhD olalekan.ut PhD	Institute of Public Health, Faculty of Health Sciences Warwick Medical School	Jagiellonian University Medical College University of Warwick	Kraków Coventry	Poland United_Kingdom	Faculty of Health Sciences	Wroclaw Medical University	Wroclaw	Poland
Prof. Mr. Mr.	Stein Emil Fiseha Tolassa		Vollset Wadilo Wakayo	vollset@ui DrPH fisehawadi@ MS wakayotol@ MS	Center for Disease Burden	Norwegian Institute of Public Health Wolaita Sodo University Jimma University	Bergen Jimma	NA Ethiopia Oromia	Department of Global Public Health and Primary Care	University of Bergen	Bergen	Norway Institute fo University iSeattle WA United_Sta
Dr. Prof. Mr. Mr.	Andrea Tissa Mohsen Hassen	Hamid	Werdecker Wijeratne Yaghoubi Yimam	andrea.we PhD twi@unim@ MD mohsen.ya MA abdulhami MPH	Competence Center Mortality-Follow-Up of the German National Cohort	Federal Institute for Population Research Western Health University of Saskatchewan Mizan Tepi University	Wiesbaden Footscray Saskatoon Mizan Teferi	Hessen Victoria Saskatchewan Ethiopia		University of Melbourne	Footscray Victoria Tehran	Australia Australia Iran
Mr. Prof. Prof. Prof.	Naohiro Mustafa Maysaa	Z El Sayed	Yonemoto Younis Zaki	nyonemot@ MPH younis68@ DrPH may_s65@ PhD	Department of Biostatistics, School of Public Health	Kyoto University Jackson State University Mansoura University	Kyoto Jackson Mansoura	Japan United_States Egypt				
Dr. Prof. Prof.	Aisha Theo Mohsen	O	Jumaan Vos Naghavi	aisha@j@ PhD tvos@uw.ε PhD nagham@t PhD	Institute for Health Metrics and Evaluation Institute for Health Metrics and Evaluation Oxford Big Data Institute, Li Ka Shing Centre for Health Information and Discovery	Independent Consultant University of Washington University of Washington	Seattle Seattle Seattle	Washington United_States United_States United_States				
Prof. Prof. Prof.	Simon Christopher Ali	I. J H.	Hay Murray Mokdad	sihay@uw. DSc cjlm@uw.ε DPhil mokdaa@t PhD	Institute for Health Metrics and Evaluation	University of Washington University of Washington	Oxford Seattle Seattle	United_Kin Evaluation United_States United_States	Institute for Health Metrics and Evaluation	University of Washington	Seattle Washington	United_States